

Rabbit MEF-2 Polyclonal Antibody | anti-MEF-2 antibody

MEF-2 (phospho Thr319) Polyclonal Antibody

CATALOG # AAA293913	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---

PRODUCT SYNONYMS

MEF-2; Polyclonal Antibody; MEF-2 (phospho Thr319) Polyclonal Antibody; MEF2A; MEF2; Myocyte-specific enhancer factor 2A; Serum response factor-like protein 1; anti-MEF-2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-MEF-2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES


WB (Western Blot)


Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MEF-2 antibody

Transcriptional activator which binds specifically to the MEF2 element, 5'-YTA[AT] (4)TAR-3', found in numerous muscle-specific genes. Also involved in the activation of numerous growth factor- and stress-induced genes. Mediates cellular functions not only in skeletal and cardiac muscle development, but also in neuronal differentiation and survival. Plays diverse roles in the control of cell growth, survival and apoptosis via p38 MAPK signaling in muscle-specific and/or growth factor-related transcription. In cerebellar granule neurons, phosphorylated and sumoylated MEF2A represses transcription of NUR77 promoting synaptic differentiation. Associates with chromatin to the ZNF16 promoter.

NCBI AND UNIPROT INFORMATION

NCBI GI #	195972801
-----------	-----------

NCBI GeneID	4205
NCBI Accession #	NP_001124398.1
NCBI GenBank Nucleotide #	NM_001130926.1
UniProt Accession #	Q02078; O43814; Q14223; Q14224; Q59GX4; Q7Z6C9; Q96D14; B4DFQ7; F6XG23
Molecular Weight	54,811 Da
NCBI Official Full Name	myocyte-specific enhancer factor 2A isoform 2
NCBI Official Synonym Full Names	myocyte enhancer factor 2A
NCBI Official Symbol	MEF2A
NCBI Official Synonym Symbols	mef2; ADCAD1; RSRFC4; RSRFC9
NCBI Protein Information	myocyte-specific enhancer factor 2A; serum response factor-like protein 1; MADS box transcription enhancer factor 2, polypeptide A (myocyte enhancer factor 2A)
UniProt Protein Name	Myocyte-specific enhancer factor 2A
UniProt Gene Name	MEF2A
UniProt Synonym Gene Names	MEF2
UniProt Entry Name	MEF2A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mef-2-antibody-immunohistochemistry-western-blot-elisa/293913>