

Rabbit HDAC2 Polyclonal Antibody | anti-HDAC2 antibody

HDAC2 (phospho Ser394) Polyclonal Antibody

CATALOG # AAA293943	HOST Rabbit	REACTIVITY Human, Mouse, Rat, Monkey	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

HDAC2; Polyclonal Antibody; HDAC2 (phospho Ser394) Polyclonal Antibody; Histone deacetylase 2; HD2; anti-HDAC2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Applicable Applications for anti-HDAC2 antibody	WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-HDAC2 antibody

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Forms transcriptional repressor complexes by associating with MAD, SIN3, YY1 and N-COR. Interacts in the late S-phase of DNA-replication with DNMT1 in the other transcriptional repressor complex composed of DNMT1, DMAP1, PCNA, CAF1. Deacetylates TSHZ3 and regulates its transcriptional repressor activity. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development. May be involved in the transcriptional repression of circadian target genes, such as PER1, mediated by CRY1 through histone deacetylation. Involved in MTA1-mediated transcriptional corepression of TFF1 and CDKN1A.

NCBI AND UNIPROT INFORMATION

NCBI GI #	293336691
NCBI GeneID	3066
NCBI Accession #	NP_001518.3
NCBI GenBank Nucleotide #	NM_001527.3
UniProt Accession #	Q92769; Q5SRI8; Q5SZ86; Q8NEH4; B4DL58; E1P561
Molecular Weight	55,364 Da
NCBI Official Full Name	histone deacetylase 2
NCBI Official Synonym Full Names	histone deacetylase 2
NCBI Official Symbol	HDAC2
NCBI Official Synonym Symbols	HD2; RPD3; YAF1
NCBI Protein Information	histone deacetylase 2; YY1-associated factor 1; transcriptional regulator homolog RPD3
UniProt Protein Name	Histone deacetylase 2
UniProt Gene Name	HDAC2
UniProt Synonym Gene Names	HD2
UniProt Entry Name	HDAC2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hdac2-antibody-western-blot-elisa/293943>