

Rabbit Cyclin D1 Polyclonal Antibody | anti-CCND1 antibody

Cyclin D1 (phospho Thr286) Polyclonal Antibody

CATALOG # AAA293949	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Cyclin D1; Polyclonal Antibody; Cyclin D1 (phospho Thr286) Polyclonal Antibody; CCND1; BCL1; PRAD1; G1/S-specific cyclin-D1; B-cell lymphoma 1 protein; BCL-1; BCL-1 oncogene; PRAD1 oncogene; anti-CCND1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-CCND1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Notes	ECO:0000303 PubMed:8204893, ECO:0000312 HGNC: HGNC:1582
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CCND1 antibody	Regulatory component of the cyclin D1-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G (1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G (1) phase. Hypophosphorylates RB1 in early G (1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity. Component of the ternary complex, cyclin D1/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex. Exhibits transcriptional corepressor activity with INSM1 on the NEUROD1 and INS promoters in a cell cycle-independent manner.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	16950655
NCBI GeneID	595
NCBI Accession #	NP_444284.1
NCBI GenBank Nucleotide #	NM_053056.2
UniProt Accession #	P24385; Q6LEF0
Molecular Weight	33,729 Da
NCBI Official Full Name	G1/S-specific cyclin-D1
NCBI Official Synonym Full Names	cyclin D1
NCBI Official Symbol	CCND1
NCBI Official Synonym Symbols	BCL1; PRAD1; U21B31; D11S287E
NCBI Protein Information	G1/S-specific cyclin-D1; BCL-1 oncogene; PRAD1 oncogene; B-cell CLL/lymphoma 1; B-cell lymphoma 1 protein
UniProt Protein Name	G1/S-specific cyclin-D1
UniProt Gene Name	CCND1
UniProt Synonym Gene Names	BCL1; PRAD1; BCL-1
UniProt Entry Name	CCND1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cyclin-d1-antibody-immunohistochemistry-western-blot-elisa/293949>