

Rabbit PLC gamma1 Polyclonal Antibody | anti-PLCgamma1 antibody

PLC gamma1 (phospho Tyr771) Polyclonal Antibody

CATALOG # AAA294034	HOST Rabbit	REACTIVITY Human, Mouse, Rat, Monkey	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PLC gamma1; Polyclonal Antibody; PLC gamma1 (phospho Tyr771) Polyclonal Antibody; PLCG1; PLC1; 1-phosphatidylinositol 4; 5-bisphosphate phosphodiesterase gamma-1; PLC-148; Phosphoinositide phospholipase C-gamma-1; Phospholipase C-II; PLC-II; Phospholipase C-gamma-1; PLC-gamma-1; anti-PLCgamma1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Applicable Applications for anti-PLCgamma1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Cellular Location	Cell projection, lamellipodium. Cell projection, ruffle. Note=Rapidly redistributed to ruffles and lamellipodia structures in response to epidermal growth factor (EGF) treatment.
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



IHC (Immunohiostchemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-PLCgamma1 antibody

Mediates the production of the second messenger molecules diacylglycerol (DAG) and inositol 1, 4, 5-trisphosphate (IP3). Plays an important role in the regulation of intracellular signaling cascades. Becomes activated in response to ligand- mediated activation of receptor-type tyrosine kinases, such as PDGFRA, PDGFRB, FGFR1, FGFR2, FGFR3 and FGFR4. Plays a role in actin reorganization and cell migration.

NCBI AND UNIPROT INFORMATION

NCBI GI #	33598948
NCBI GeneID	5335
NCBI Accession #	NP_002651.2
NCBI GenBank Nucleotide #	NM_002660.2
UniProt Accession #	P19174; Q2V575; B7ZLY7; B9EGH4; E1P5W4
Molecular Weight	148,532 Da
NCBI Official Full Name	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1 isoform a
NCBI Official Synonym Full Names	phospholipase C, gamma 1
NCBI Official Symbol	PLCG1
NCBI Official Synonym Symbols	PLC1; NCKAP3; PLC-II; PLC148; PLCgamma1
NCBI Protein Information	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1; PLC-148; PLC-gamma-1; phospholipase C-II; phospholipase C-148; phosphoinositidase C; phospholipase C-gamma-1; inositoltrisphosphohydrolase; phosphoinositide phospholipase C; phosphatidylin
UniProt Protein Name	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1
UniProt Gene Name	PLCG1
UniProt Synonym Gene Names	PLC1; PLC-II; PLC-gamma-1
UniProt Entry Name	PLCG1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/plc-gamma1-antibody-immunohistochemistry-western-blot-elisa/294034>