

Rabbit anti-Human, Mouse Emt Polyclonal Antibody | anti-Emt antibody

Emt (phospho Tyr512) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294124	Rabbit	Human, Mouse	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Emt; Polyclonal Antibody; Emt (phospho Tyr512) Polyclonal Antibody; ITK; EMT; LYK; Tyrosine-protein kinase ITK/TSK; Interleukin-2-inducible T-cell kinase; IL-2-inducible T-cell kinase; Kinase EMT; T-cell-specific kinase; Tyrosine-protein kinase Lyk; anti-Emt antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Applicable Applications for anti-Emt antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Dilution: Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.



IHC (Immunohistochemistry)

Dilution: Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-Emt antibody	Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylation lead to the recruitment of ITK to the cell membrane, in the vicinity of the stimulated TCR receptor, where it is phosphorylated by LCK. Phosphorylation leads to ITK autophosphorylation and full activation. Once activated, phosphorylates PLCG1, leading to the activation of this lipase and subsequent cleavage of its substrates. In turn, the endoplasmic reticulum releases calcium in the cytoplasm and the nuclear activator of activated T-cells (NFAT) translocates into the nucleus to perform its transcriptional duty. Phosphorylates 2 essential adapter proteins: the linker for activation of T-cells/LAT protein and LCP2. Then, a large number of signaling molecules such as VAV1 are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation (PubMed:12682224, PubMed:21725281). Phosphorylates TBX21 at 'Tyr-530' and mediates its interaction with GATA3 (By similarity).
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	585361
NCBI GeneID	3702
UniProt Accession #	Q08881; Q32ML7; B2R752
Molecular Weight	620
NCBI Official Full Name	Tyrosine-protein kinase ITK/TSK
NCBI Official Synonym Full Names	IL2-inducible T-cell kinase
NCBI Official Symbol	ITK
NCBI Official Synonym Symbols	EMT; LYK; LPFS1; PSCTK2
NCBI Protein Information	tyrosine-protein kinase ITK/TSK; kinase EMT; T-cell-specific kinase; tyrosine-protein kinase LYK; IL-2-inducible T cell kinase; IL-2-inducible T-cell kinase; homolog of mouse T-cell itk/tsk; interleukin-2-inducible T cell kinase; interleukin-2-inducible T
UniProt Protein Name	Tyrosine-protein kinase ITK/TSK
UniProt Gene Name	ITK
UniProt Synonym Gene Names	EMT; LYK; IL-2-inducible T-cell kinase
UniProt Entry Name	ITK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/emt-antibody-immunohistochemistry-western-blot-elisa/294124>