

Rabbit EDG-1 Polyclonal Antibody | anti-EDG-1 antibody

EDG-1 (phospho Thr236) Polyclonal Antibody

CATALOG # AAA294142	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

EDG-1; Polyclonal Antibody; EDG-1 (phospho Thr236) Polyclonal Antibody; S1PR1; CHEDG1; EDG1; Sphingosine 1-phosphate receptor 1; S1P receptor 1; S1P1; Endothelial differentiation G-protein coupled receptor 1; Sphingosine 1-phosphate receptor Edg-1; S1P receptor Edg-1; CD antigen CD363; anti-EDG-1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-EDG-1 antibody	IF (Immunofluorescence), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.



IF (Immunofluorescence)

Dilution: Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-EDG-1 antibody	G-protein coupled receptor for the bioactive lysosphingolipid sphingosine 1-phosphate (S1P) that seems to be coupled to the G (i) subclass of heteromeric G proteins. Signaling leads to the activation of RAC1, SRC, PTK2/FAK1 and MAP kinases. Plays an important role in cell migration, probably via its role in the reorganization of the actin cytoskeleton and the formation of lamellipodia in response to stimuli that increase the activity of the sphingosine kinase SPHK1. Required for normal chemotaxis toward sphingosine 1-phosphate. Required for normal embryonic heart development and normal cardiac morphogenesis. Plays an important role in the regulation of sprouting angiogenesis and vascular maturation. Inhibits sprouting angiogenesis to prevent excessive sprouting during blood vessel development. Required for normal egress of mature T-cells from the thymus into the blood stream and into peripheral lymphoid organs. Plays a role in the migration of osteoclast precursor cells, the regulation of bone mineralization and bone homeostasis (By similarity). Plays a role in responses to oxidized 1-palmitoyl-2-arachidonoyl-sn-glycero-3- phosphocholine by pulmonary endothelial cells and in the protection against ventilator-induced lung injury.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	13027636
NCBI GeneID	1901
NCBI Accession #	NP_001391.2
NCBI GenBank Nucleotide #	NM_001400.4
UniProt Accession #	P21453; Q9BYY4; Q9NYN8; D3DT66
Molecular Weight	42,811 Da
NCBI Official Full Name	sphingosine 1-phosphate receptor 1
NCBI Official Synonym Full Names	sphingosine-1-phosphate receptor 1
NCBI Official Symbol	S1PR1
NCBI Official Synonym Symbols	EDG1; S1P1; CD363; ECGF1; EDG-1; CHEDG1; D1S3362
NCBI Protein Information	sphingosine 1-phosphate receptor 1; S1P receptor 1; S1P receptor Edg-1; sphingosine 1-phosphate receptor EDG1; sphingosine 1-phosphate receptor Edg-1; endothelial differentiation G-protein coupled receptor 1; endothelial differentiation, sphingolipid G-pr
UniProt Protein Name	Sphingosine 1-phosphate receptor 1
UniProt Gene Name	S1PR1
UniProt Synonym Gene Names	CHEDG1; EDG1; S1P receptor 1; S1P1
UniProt Entry Name	S1PR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/edg-1-antibody-immunofluorescence-western-blot-elisa/294142>