

Rabbit Catenin-beta1 Polyclonal Antibody | anti-CTNNB1 antibody

Catenin-beta1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294397	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Catenin-beta1; Polyclonal Antibody; Catenin-beta1 Polyclonal Antibody; CTNNB1; CTNNB; OK/SW-cl.35; Catenin beta-1; Beta-catenin; anti-CTNNB1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-CTNNB1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CTNNB1 antibody	Key downstream component of the canonical Wnt signaling pathway. In the absence of Wnt, forms a complex with AXIN1, AXIN2, APC, CSNK1A1 and GSK3B that promotes phosphorylation on N-terminal Ser and Thr residues and ubiquitination of CTNNB1 via BTRC and its subsequent degradation by the proteasome. In the presence of Wnt ligand, CTNNB1 is not ubiquitinated and accumulates in the nucleus, where it acts as a coactivator for transcription factors of the TCF/LEF family, leading to activate Wnt responsive genes. Involved in the regulation of cell adhesion, as component of an E- cadherin: catenin adhesion complex. Acts as a negative regulator of centrosome cohesion. Involved in the CDK2/PTPN6/CTNNB1/CEACAM1 pathway of insulin internalization. Blocks anoikis of malignant kidney and intestinal epithelial cells and promotes their anchorage-independent growth by down-regulating DAPK2. Disrupts PML function and PML-NB formation by inhibiting RANBP2-mediated sumoylation of PML (PubMed:18077326, PubMed:18086858, PubMed:18957423, PubMed:21262353, PubMed:22647378, PubMed:22699938, PubMed:22155184). Promotes neurogenesis by maintaining sympathetic neuroblasts within the cell cycle (By similarity).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	148233338
NCBI GeneID	1499
NCBI Accession #	NP_001091679.1
NCBI GenBank Nucleotide #	NM_001098209.1
UniProt Accession #	P35222; Q8NEW9; Q8NI94; Q9H391; A8K1L7
Molecular Weight	85,497 Da
NCBI Official Full Name	catenin beta-1
NCBI Official Synonym Full Names	catenin (cadherin-associated protein), beta 1, 88kDa
NCBI Official Symbol	CTNNB1
NCBI Official Synonym Symbols	CTNNB; MRD19; armadillo
NCBI Protein Information	catenin beta-1
UniProt Protein Name	Catenin beta-1
UniProt Gene Name	CTNNB1
UniProt Synonym Gene Names	CTNNB
UniProt Entry Name	CTNB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.