

Rabbit anti-Human CD3-delta Polyclonal Antibody | anti-CD3-delta antibody

CD3-delta Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294429	Rabbit	Human	Western Blot


PRODUCT SYNONYMS

CD3-delta; Polyclonal Antibody; CD3-delta Polyclonal Antibody; CD3D; T3D; T-cell surface glycoprotein CD3 delta chain; T-cell receptor T3 delta chain; CD antigen CD3d; anti-CD3-delta antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD3-delta antibody	WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CD3-delta antibody

Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:2470098). In addition of this role of signal transduction in T-cell activation, CD3D plays an essential role in thymocyte differentiation. Indeed, participates in correct intracellular TCR-CD3 complex assembly and surface expression. In absence of a functional TCR-CD3 complex, thymocytes are unable to differentiate properly. Interacts with CD4 and CD8 and thus serves to establish a functional link between the TCR and coreceptors CD4 and CD8, which is needed for activation and positive selection of CD4 or CD8 T-cells (PubMed:12215456).

NCBI AND UNIPROT INFORMATION

NCBI GI #	115985
NCBI GeneID	915
UniProt Accession #	P04234; A8MVP6
Molecular Weight	171
NCBI Official Full Name	T-cell surface glycoprotein CD3 delta chain
NCBI Official Synonym Full Names	CD3d molecule, delta (CD3-TCR complex)
NCBI Official Symbol	CD3D
NCBI Official Synonym Symbols	T3D; IMD19; CD3-DELTA
NCBI Protein Information	T-cell surface glycoprotein CD3 delta chain; OKT3, delta chain; CD3 antigen, delta subunit; T-cell receptor T3 delta chain; CD3d antigen, delta polypeptide (TIT3 complex)
UniProt Protein Name	T-cell surface glycoprotein CD3 delta chain
UniProt Gene Name	CD3D
UniProt Synonym Gene Names	T3D
UniProt Entry Name	CD3D_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd3-delta-antibody-western-blot-elisa/294429>