

Rabbit Cdc20 Polyclonal Antibody | anti-Cdc20 antibody

Cdc20 Polyclonal Antibody

CATALOG # AAA294440	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Cdc20; Polyclonal Antibody; Cdc20 Polyclonal Antibody; CDC20; Cell division cycle protein 20 homolog; p55CDC; anti-Cdc20 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-Cdc20 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.

Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-Cdc20 antibody

Required for full ubiquitin ligase activity of the anaphase promoting complex/cyclosome (APC/C) and may confer substrate specificity upon the complex. Is regulated by MAD2L1: in metaphase the MAD2L1-CDC20-APC/C ternary complex is inactive and in anaphase the CDC20-APC/C binary complex is active in degrading substrates. The CDC20-APC/C complex positively regulates the formation of synaptic vesicle clustering at active zone to the presynaptic membrane in postmitotic neurons. CDC20-APC/C-induced degradation of NEUROD2 induces presynaptic differentiation.

NCBI AND UNIPROT INFORMATION

NCBI GI #	118402582
NCBI GeneID	991
NCBI Accession #	NP_001246.2
NCBI GenBank Nucleotide #	NM_001255.2
UniProt Accession #	Q12834; Q5JUY4; Q9BW56; Q9UQI9; B2R6Z6; D3DPJ1
Molecular Weight	54,723 Da
NCBI Official Full Name	cell division cycle protein 20 homolog
NCBI Official Synonym Full Names	cell division cycle 20
NCBI Official Symbol	CDC20
NCBI Official Synonym Symbols	CDC20A; p55CDC; bA276H19.3
NCBI Protein Information	cell division cycle protein 20 homolog; cell division cycle 20 homolog; CDC20 cell division cycle 20 homolog
UniProt Protein Name	Cell division cycle protein 20 homolog
UniProt Gene Name	CDC20
UniProt Entry Name	CDC20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cdc20-antibody-immunohistochemistry-western-blot-elisa/294440>