

Rabbit anti-Human c-Fms Polyclonal Antibody | anti-c-Fms antibody

c-Fms Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294473	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

c-Fms; Polyclonal Antibody; c-Fms Polyclonal Antibody; CSF1R; FMS; Macrophage colony-stimulating factor 1 receptor; CSF-1 receptor; CSF-1-R; CSF-1R; M-CSF-R; Proto-oncogene c-Fms; CD antigen CD115; anti-c-Fms antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-c-Fms antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-c-Fms antibody

Tyrosine-protein kinase that acts as cell-surface receptor for CSF1 and IL34 and plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes, such as macrophages and monocytes. Promotes the release of proinflammatory chemokines in response to IL34 and CSF1, and thereby plays an important role in innate immunity and in inflammatory processes. Plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone and tooth development. Required for normal male and female fertility, and for normal development of milk ducts and acinar structures in the mammary gland during pregnancy. Promotes reorganization of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration, and promotes cancer cell invasion. Activates several signaling pathways in response to ligand binding. Phosphorylates PIK3R1, PLCG2, GRB2, SLA2 and CBL. Activation of PLCG2 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1, 4, 5- trisphosphate, that then lead to the activation of protein kinase C family members, especially PRKCD. Phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, leads to activation of the AKT1 signaling pathway. Activated CSF1R also mediates activation of the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1, and of the SRC family kinases SRC, FYN and YES1. Activated CSF1R transmits signals both via proteins that directly interact with phosphorylated tyrosine residues in its intracellular domain, or via adapter proteins, such as GRB2. Promotes activation of STAT family members STAT3, STAT5A and/or STAT5B. Promotes tyrosine phosphorylation of SHC1 and INPP5D/SHIP- 1. Receptor signaling is down-regulated by protein phosphatases, such as INPP5D/SHIP-1, that dephosphorylate the receptor and its downstream effectors, and by rapid internalization of the activated receptor.

NCBI AND UNIPROT INFORMATION

NCBI GI #	27262659
NCBI GeneID	1436
NCBI Accession #	NP_005202.2
NCBI GenBank Nucleotide #	NM_005211.3
UniProt Accession #	P07333; Q6LDW5; Q6LDY4; Q86VW7; D3DQG2
Molecular Weight	107,984 Da
NCBI Official Full Name	macrophage colony-stimulating factor 1 receptor
NCBI Official Synonym Full Names	colony stimulating factor 1 receptor
NCBI Official Symbol	CSF1R
NCBI Official Synonym Symbols	FMS; CSFR; FIM2; HDLS; C-FMS; CD115; CSF-1R; M-CSF-R
NCBI Protein Information	macrophage colony-stimulating factor 1 receptor; CD115 antigen; CSF-1 receptor; FMS proto-oncogene; proto-oncogene c-Fms; macrophage colony stimulating factor I receptor; McDonough feline sarcoma viral (v-fms) oncogene homolog
UniProt Protein Name	Macrophage colony-stimulating factor 1 receptor
UniProt Gene Name	CSF1R
UniProt Synonym Gene Names	FMS; CSF-1-R; CSF-1R; M-CSF-R
UniProt Entry Name	CSF1R_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/c-fms-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/294473>