

Rabbit COT Polyclonal Antibody | anti-COT antibody

COT Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294543	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

COT; Polyclonal Antibody; COT Polyclonal Antibody; MAP3K8; ESTF; Mitogen-activated protein kinase kinase kinase 8; Cancer Osaka thyroid oncogene; Proto-oncogene c-Cot; Serine/threonine-protein kinase cot; Tumor progression locus 2; TPL-2; anti-COT antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-COT antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-COT antibody	Required for lipopolysaccharide (LPS)-induced, TLR4- mediated activation of the MAPK/ERK pathway in macrophages, thus being critical for production of the proinflammatory cytokine TNF- alpha (TNF) during immune responses. Involved in the regulation of T-helper cell differentiation and IFNG expression in T-cells. Involved in mediating host resistance to bacterial infection through negative regulation of type I interferon (IFN) production. In vitro, activates MAPK/ERK pathway in response to IL1 in an IRAK1-independent manner, leading to up-regulation of IL8 and CCL4. Transduces CD40 and TNFRSF1A signals that activate ERK in B- cells and macrophages, and thus may play a role in the regulation of immunoglobulin production. May also play a role in the transduction of TNF signals that activate JNK and NF-kappa-B in some cell types. In adipocytes, activates MAPK/ERK pathway in an IKBKB-dependent manner in response to IL1B and TNF, but not insulin, leading to induction of lipolysis. Plays a role in the cell cycle. Isoform 1 shows some transforming activity, although it is much weaker than that of the activated oncogenic variant.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	346644795
NCBI GeneID	1326
NCBI Accession #	NP_001231063.1
NCBI GenBank Nucleotide #	NM_001244134.1
UniProt Accession #	P41279; Q14275; Q5T855; Q9HC81; A8K2Q5; D3DRX1
Molecular Weight	52,925 Da
NCBI Official Full Name	mitogen-activated protein kinase kinase kinase 8
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase kinase 8
NCBI Official Symbol	MAP3K8
NCBI Official Synonym Symbols	COT; EST; ESTF; TPL2; MEKK8; Tpl-2; c-COT
NCBI Protein Information	mitogen-activated protein kinase kinase kinase 8; proto-oncogene c-Cot; tumor progression locus 2; Ewing sarcoma transformant; cot (cancer Osaka thyroid) oncogene; proto-oncogene serine/threonine protein kinase
UniProt Protein Name	Mitogen-activated protein kinase kinase kinase 8
UniProt Gene Name	MAP3K8
UniProt Synonym Gene Names	COT; ESTF; TPL-2
UniProt Entry Name	M3K8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cot-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/294543>