

Rabbit anti-Human, Mouse CREB-2 Polyclonal Antibody | anti-CREB-2 antibody

CREB-2 Polyclonal Antibody

CATALOG # AAA294553	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Western Blot
-------------------------------	-----------------------	-----------------------------------	-------------------------------------


PRODUCT SYNONYMS

CREB-2; Polyclonal Antibody; CREB-2 Polyclonal Antibody; ATF4; CREB2; TXREB; Cyclic AMP-dependent transcription factor ATF-4; cAMP-dependent transcription factor ATF-4; Activating transcription factor 4; Cyclic AMP-responsive element-binding protein 2; cAMP-responsive element-binding prot; anti-CREB-2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Applicable Applications for anti-CREB-2 antibody	WB (Western Blot)
Notes	ECO:0000303 PubMed:2516827, ECO:0000312 HGNC: HGNC:786
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CREB-2 antibody	Transcriptional activator. Binds the cAMP response element (CRE)(consensus:5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Cooperates with FOXO1 in osteoblasts to regulate glucose homeostasis through suppression of beta-cell production and decrease in insulin production (By similarity). It binds to a Tax-responsive enhancer element in the long terminal repeat of HTLV-I. Regulates the induction of DDIT3/CHOP and asparagine synthetase (ASNS) in response to endoplasmic reticulum (ER) stress. In concert with DDIT3/CHOP, activates the transcription of TRIB3 and promotes ER stress-induced neuronal apoptosis by regulating the transcriptional induction of BBC3/PUMA. Activates transcription of SIRT4. Regulates the circadian expression of the core clock component PER2 and the serotonin transporter SLC6A4. Binds in a circadian time-dependent manner to the cAMP response elements (CRE) in the SLC6A4 and PER2 promoters and periodically activates the transcription of these genes. During ER stress response, activates the transcription of NLRP1, possibly in concert with other factors (PubMed:26086088).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	33469976
NCBI GeneID	468
NCBI Accession #	NP_001666.2
NCBI GenBank Nucleotide #	NM_001675.2
UniProt Accession #	P18848; Q9UH31
Molecular Weight	40 kDa
NCBI Official Full Name	cyclic AMP-dependent transcription factor ATF-4
NCBI Official Synonym Full Names	activating transcription factor 4
NCBI Official Symbol	ATF4
NCBI Official Synonym Symbols	CREB2; TXREB; CREB-2; TAXREB67
NCBI Protein Information	cyclic AMP-dependent transcription factor ATF-4; DNA-binding protein TAXREB67; tax-responsive enhancer element B67; cAMP response element-binding protein 2; cAMP-dependent transcription factor ATF-4; cAMP-responsive element-binding protein 2; cyclic AMP-r
UniProt Protein Name	Cyclic AMP-dependent transcription factor ATF-4
UniProt Gene Name	ATF4
UniProt Synonym Gene Names	CREB2; TXREB; cAMP-dependent transcription factor ATF-4; CREB-2; cAMP-responsive element-binding protein 2; TaxREB67
UniProt Entry Name	ATF4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.