

Rabbit Desmin Polyclonal Antibody | anti-DES antibody

Desmin Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294665	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Desmin; Polyclonal Antibody; Desmin Polyclonal Antibody; DES; anti-DES antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-DES antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Cellular Location	Cytoplasm, myofibril, sarcomere, Z line. Cytoplasm. Cell membrane, sarcolemma. Nucleus Note=Localizes in the intercalated disks which occur at the Z line of cardiomyocytes (PubMed:24200904, PubMed:26724190). Localizes in the nucleus exclusively in differentiating cardiac progenitor cells and premature cardiomyocytes (By similarity)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-DES antibody

Muscle-specific type III intermediate filament essential for proper muscular structure and function. Plays a crucial role in maintaining the structure of sarcomeres, inter-connecting the Z-disks and forming the myofibrils, linking them not only to the sarcolemmal cytoskeleton, but also to the nucleus and mitochondria, thus providing strength for the muscle fiber during activity (PubMed:25394388, PubMed:26724190). May act as a sarcomeric microtubule-anchoring protein: specifically associates with detyrosinated tubulin-alpha chains, leading to buckled microtubules and mechanical resistance to contraction. Contributes to the transcriptional regulation of the NKX2-5 gene in cardiac progenitor cells during a short period of cardiomyogenesis and in cardiac side population stem cells in the adult. Plays a role in maintaining an optimal conformation of nebulin (NEB) on heart muscle sarcomeres to bind and recruit cardiac alpha-actin (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	55749932
NCBI GeneID	1674
NCBI Accession #	NP_001918.3
NCBI GenBank Nucleotide #	NM_001927.3
UniProt Accession #	P17661; Q15787; Q549R7; Q549R8; Q549R9; Q8IZR1; Q8IZR6; Q8NES2; Q8NEU6; Q8TAC4; Q8TCX2; Q8TD99
Molecular Weight	53,536 Da
NCBI Official Full Name	desmin
NCBI Official Synonym Full Names	desmin
NCBI Official Symbol	DES
NCBI Official Synonym Symbols	CSM1; CSM2
NCBI Protein Information	desmin; intermediate filament protein
UniProt Protein Name	Desmin
UniProt Gene Name	DES
UniProt Entry Name	DESM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.