

Rabbit GluR-1 Polyclonal Antibody | anti-GluR-1 antibody

GluR-1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294909	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

GluR-1; Polyclonal Antibody; GluR-1 Polyclonal Antibody; GRIA1; GLUH1; GLUR1; Glutamate receptor 1; AMPA-selective glutamate receptor 1; GluR-A; GluR-K1; Glutamate receptor ionotropic; AMPA 1; GluA1; anti-GluR-1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-GluR-1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES


WB (Western Blot)


Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-GluR-1 antibody

Ionotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L- glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of glutamate.

NCBI AND UNIPROT INFORMATION

NCBI GI #	167001419
-----------	-----------

NCBI GeneID	2890
NCBI Accession #	NP_000818.2
NCBI GenBank Nucleotide #	NM_000827.3
UniProt Accession #	P42261; Q2NKM6; B7Z2S0; B7Z3F6; D3DQI4
Molecular Weight	101,506 Da
NCBI Official Full Name	glutamate receptor 1 isoform 1
NCBI Official Synonym Full Names	glutamate receptor, ionotropic, AMPA 1
NCBI Official Symbol	GRIA1
NCBI Official Synonym Symbols	GLUH1; GLUR1; GLURA; GluA1; HBGR1
NCBI Protein Information	glutamate receptor 1; AMPA 1; gluR-1; gluR-A; gluR-K1; AMPA-selective glutamate receptor 1
UniProt Protein Name	Glutamate receptor 1
UniProt Gene Name	GRIA1
UniProt Synonym Gene Names	GLUH1; GLUR1; GluR-1; GluA1
UniProt Entry Name	GRIA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/gluR-1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/294909>