

Rabbit GluR-2 Polyclonal Antibody | anti-GluR-2 antibody

GluR-2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294911	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

GluR-2; Polyclonal Antibody; GluR-2 Polyclonal Antibody; GRIA2; GLUR2; Glutamate receptor 2; AMPA-selective glutamate receptor 2; GluR-B; GluR-K2; Glutamate receptor ionotropic; AMPA 2; GluA2; anti-GluR-2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-GluR-2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-GluR-2 antibody	Receptor for glutamate that functions as ligand-gated ion channel in the central nervous system and plays an important role in excitatory synaptic transmission. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L- glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of glutamate. Through complex formation with NSG1, GRIP1 and STX12 controls the intracellular fate of AMPAR and the endosomal sorting of the GRIA2 subunit toward recycling and membrane targeting (By similarity).
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	74275354
NCBI GeneID	2891
NCBI Accession #	NP_000817.2
NCBI GenBank Nucleotide #	NM_000826.3

UniProt Accession #	P42262; Q96FP6
Molecular Weight	98,821 Da
NCBI Official Full Name	glutamate receptor 2 isoform 1
NCBI Official Synonym Full Names	glutamate receptor, ionotropic, AMPA 2
NCBI Official Symbol	GRIA2
NCBI Official Synonym Symbols	GLUR2; GLURB; GluA2; HBGR2; GluR-K2
NCBI Protein Information	glutamate receptor 2; glur-2; glur-B; AMPA-selective glutamate receptor 2
UniProt Protein Name	Glutamate receptor 2
UniProt Gene Name	GRIA2
UniProt Synonym Gene Names	GLUR2; GluR-2; GluA2
UniProt Entry Name	GRIA2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/glur-2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/294911>