

Rabbit HDAC3 Polyclonal Antibody | anti-HDAC3 antibody

HDAC3 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294978	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS


HDAC3; Polyclonal Antibody; HDAC3 Polyclonal Antibody; Histone deacetylase 3; HD3; RPD3-2; SMAP45; anti-HDAC3 antibody


PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-HDAC3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-HDAC3 antibody

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates. Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys- 27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1; increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor. Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation (PubMed:23911289). Contributes, together with XBP1 isoform 1, to the activation of NFE2L2-mediated HMOX1 transcription factor gene expression in a PI (3)K/mTORC2/Akt-dependent signaling pathway leading to endothelial cell (EC) survival under disturbed flow/oxidative stress (PubMed:25190803).

NCBI AND UNIPROT INFORMATION

NCBI GI #	13128862
NCBI GeneID	8841
NCBI Accession #	NP_003874.2
NCBI GenBank Nucleotide #	NM_003883.3
UniProt Accession #	O15379; O43268; Q9UEI5; Q9UEV0; D3DQE1
Molecular Weight	48,848 Da
NCBI Official Full Name	histone deacetylase 3
NCBI Official Synonym Full Names	histone deacetylase 3
NCBI Official Symbol	HDAC3
NCBI Official Synonym Symbols	HD3; RPD3; RPD3-2
NCBI Protein Information	histone deacetylase 3; SMAP45
UniProt Protein Name	Histone deacetylase 3
UniProt Gene Name	HDAC3
UniProt Synonym Gene Names	HD3
UniProt Entry Name	HDAC3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hdac3-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/294978>