

Rabbit anti-Human, Mouse JAB1 Polyclonal Antibody | anti-JAB1 antibody

JAB1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA295102	Rabbit	Human, Mouse	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

JAB1; Polyclonal Antibody; JAB1 Polyclonal Antibody; COPS5; CSN5; COP9 signalosome complex subunit 5; SGN5; Signalosome subunit 5; Jun activation domain-binding protein 1; anti-JAB1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Applicable Applications for anti-JAB1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000.
Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-JAB1 antibody

Probable protease subunit of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of the SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1 and IRF8, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. In the complex, it probably acts as the catalytic center that mediates the cleavage of Nedd8 from cullins. It however has no metalloprotease activity by itself and requires the other subunits of the CSN complex. Interacts directly with a large number of proteins that are regulated by the CSN complex, confirming a key role in the complex. Promotes the proteasomal degradation of BRSK2.

NCBI AND UNIPROT INFORMATION

NCBI GI #	38027923
NCBI GeneID	10987
NCBI Accession #	NP_006828.2
NCBI GenBank Nucleotide #	NM_006837.2
UniProt Accession #	Q92905; O15386; Q6AW95; Q86WQ4; Q9BQ17
Molecular Weight	37,579 Da
NCBI Official Full Name	COP9 signalosome complex subunit 5
NCBI Official Synonym Full Names	COP9 signalosome subunit 5
NCBI Official Symbol	COPS5
NCBI Official Synonym Symbols	CSN5; JAB1; SGN5; MOV-34
NCBI Protein Information	COP9 signalosome complex subunit 5; 38 kDa Mov34 homolog; signalosome subunit 5; jun activation domain-binding protein 1; COP9 constitutive photomorphogenic homolog subunit 5
UniProt Protein Name	COP9 signalosome complex subunit 5
UniProt Gene Name	COPS5
UniProt Synonym Gene Names	CSN5; JAB1; SGN5
UniProt Entry Name	CSN5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/jab1-antibody-immunohistochemistry-western-blot-elisa/295102>