

Rabbit anti-Human, Mouse MDA5 Polyclonal Antibody | anti-MDA5 antibody

MDA5 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA295204	Rabbit	Human, Mouse	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MDA5; Polyclonal Antibody; MDA5 Polyclonal Antibody; IFIH1; RH116; Interferon-induced helicase C domain-containing protein 1; Clinically amyopathic dermatomyositis autoantigen 140 kDa; CADM-140 autoantigen; Helicase with 2 CARD domains; Helicard; Interferon-induced with helicase C domain; anti-MDA5 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Applicable Applications for anti-MDA5 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MDA5 antibody

Innate immune receptor which acts as a cytoplasmic sensor of viral nucleic acids and plays a major role in sensing viral infection and in the activation of a cascade of antiviral responses including the induction of type I interferons and proinflammatory cytokines. Its ligands include mRNA lacking 2'-O- methylation at their 5' cap and long-dsRNA (>1 kb in length). Upon ligand binding it associates with mitochondria antiviral signaling protein (MAVS/IPS1) which activates the IKK-related kinases: TBK1 and IKKε which phosphorylate interferon regulatory factors: IRF3 and IRF7 which in turn activate transcription of antiviral immunological genes, including interferons (IFNs); IFN-α and IFN-β. Responsible for detecting the Picornaviridae family members such as encephalomyocarditis virus (EMCV) and mengo encephalomyocarditis virus (ENMG). Can also detect other viruses such as dengue virus (DENV), west Nile virus (WNV), and reovirus. Also involved in antiviral signaling in response to viruses containing a dsDNA genome, such as vaccinia virus. Plays an important role in amplifying innate immune signaling through recognition of RNA metabolites that are produced during virus infection by ribonuclease L (RNase L). May play an important role in enhancing natural killer cell function and may be involved in growth inhibition and apoptosis in several tumor cell lines.

NCBI AND UNIPROT INFORMATION

NCBI GI #	27886568
NCBI GeneID	64135
NCBI Accession #	NP_071451.2
NCBI GenBank Nucleotide #	NM_022168.3
UniProt Accession #	Q9BYX4; Q2NKL6; Q6DC96; Q86X56; Q96MX8; Q9H3G6
NCBI Official Full Name	interferon-induced helicase C domain-containing protein 1
NCBI Official Synonym Full Names	interferon induced with helicase C domain 1
NCBI Official Symbol	IFIH1
NCBI Official Synonym Symbols	HLcd; MDA5; MDA-5; RLR-2; IDDM19
NCBI Protein Information	interferon-induced helicase C domain-containing protein 1; helicard; CADM-140 autoantigen; RIG-I-like receptor 2; helicase with 2 CARD domains; RNA helicase-DEAD box protein 116; murabutide down-regulated protein; DEAD/H (Asp-Glu-Ala-Asp/His) box polypept
UniProt Protein Name	Interferon-induced helicase C domain-containing protein 1
UniProt Gene Name	IFIH1
UniProt Synonym Gene Names	MDA5; RH116; CADM-140 autoantigen; Helicard; MDA-5; RLR-2
UniProt Entry Name	IFIH1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mda5-antibody-immunohistochemistry-western-blot-elisa/295204>