

Rabbit MLH1 Polyclonal Antibody | anti-MLH1 antibody

MLH1 Polyclonal Antibody

CATALOG # AAA295238	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS


MLH1; Polyclonal Antibody; MLH1 Polyclonal Antibody; COCA2; DNA mismatch repair protein Mlh1; MutL protein homolog 1; anti-MLH1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-MLH1 antibody	IF (Immunofluorescence), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000.
Not yet tested in other applications.

WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MLH1 antibody

Heterodimerizes with PMS2 to form MutL alpha, a component of the post-replicative DNA mismatch repair system (MMR). DNA repair is initiated by MutS alpha (MSH2-MSH6) or MutS beta (MSH2-MSH3) binding to a dsDNA mismatch, then MutL alpha is recruited to the heteroduplex. Assembly of the MutL-MutS- heteroduplex ternary complex in presence of RFC and PCNA is sufficient to activate endonuclease activity of PMS2. It introduces single-strand breaks near the mismatch and thus generates new entry points for the exonuclease EXO1 to degrade the strand containing the mismatch. DNA methylation would prevent cleavage and therefore assure that only the newly mutated DNA strand is going to be corrected. MutL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. Also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. Heterodimerizes with MLH3 to form MutL gamma which plays a role in meiosis.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557757
NCBI GeneID	4292
NCBI Accession #	NP_000240.1
NCBI GenBank Nucleotide #	NM_000249.3
UniProt Accession #	P40692; B4DI13
Molecular Weight	84,601 Da
NCBI Official Full Name	DNA mismatch repair protein Mlh1 isoform 1
NCBI Official Synonym Full Names	mutL homolog 1, colon cancer, nonpolyposis type 2 (E. coli)
NCBI Official Symbol	MLH1
NCBI Official Synonym Symbols	FCC2; COCA2; HNPCC; hMLH1; HNPCC2
NCBI Protein Information	DNA mismatch repair protein Mlh1
UniProt Protein Name	DNA mismatch repair protein Mlh1
UniProt Gene Name	MLH1
UniProt Synonym Gene Names	COCA2
UniProt Entry Name	MLH1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mlh1-antibody-immunofluorescence-western-blot-elisa/295238>