

Rabbit Nucleophosmin Polyclonal Antibody | anti-NPM1 antibody

Nucleophosmin Polyclonal Antibody

CATALOG # AAA295418	HOST Rabbit	REACTIVITY Human, Mouse, Rat, Monkey	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Nucleophosmin; Polyclonal Antibody; Nucleophosmin Polyclonal Antibody; NPM1; NPM; Nucleolar phosphoprotein B23; Nucleolar protein NO38; Numatrin; anti-NPM1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Applicable Applications for anti-NPM1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-NPM1 antibody	Involved in diverse cellular processes such as ribosome biogenesis, centrosome duplication, protein chaperoning, histone assembly, cell proliferation, and regulation of tumor suppressors p53/TP53 and ARF. Binds ribosome presumably to drive ribosome nuclear export. Associated with nucleolar ribonucleoprotein structures and bind single-stranded nucleic acids. Acts as a chaperonin for the core histones H3, H2B and H4. Stimulates APEX1 endonuclease activity on apurinic/apyrimidinic (AP) double- stranded DNA but inhibits APEX1 endonuclease activity on AP single-stranded RNA. May exert a control of APEX1 endonuclease activity within nucleoli devoted to repair AP on rDNA and the removal of oxidized rRNA molecules. In concert with BRCA2, regulates centrosome duplication. Regulates centriole duplication: phosphorylation by PLK2 is able to trigger centriole replication. Negatively regulates the activation of EIF2AK2/PKR and suppresses apoptosis through inhibition of EIF2AK2/PKR autophosphorylation. Antagonizes the inhibitory effect of ATF5 on cell proliferation and relieves ATF5-induced G2/M blockade (PubMed:22528486). In complex with MYC enhances the transcription of MYC target genes (PubMed:25956029).
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	83641870
NCBI GeneID	4869
NCBI Accession #	NP_001032827.1
NCBI GenBank Nucleotide #	NM_001037738.2
UniProt Accession #	P06748; P08693; Q12826; Q13440; Q13441; Q14115; Q5EU94; Q5EU95; Q5EU96; A8K3N7; B5BU00; D3DQL6
Molecular Weight	32,575 Da
NCBI Official Full Name	nucleophosmin isoform 3
NCBI Official Synonym Full Names	nucleophosmin (nucleolar phosphoprotein B23, numatrin)
NCBI Official Symbol	NPM1
NCBI Official Synonym Symbols	B23; NPM
NCBI Protein Information	nucleophosmin; nucleolar protein NO38; nucleophosmin/nucleoplasmin family, member 1
UniProt Protein Name	Nucleophosmin
UniProt Gene Name	NPM1
UniProt Synonym Gene Names	NPM; NPM
UniProt Entry Name	NPM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nucleophosmin-antibody-immunohistochemistry-western-blot-elisa/295418>