

Rabbit anti-Human p53 Polyclonal Antibody | anti-p53 antibody

p53 Polyclonal Antibody

CATALOG # AAA295541	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot
-------------------------------	-----------------------	----------------------------	-------------------------------------


PRODUCT SYNONYMS

p53; Polyclonal Antibody; p53 Polyclonal Antibody; TP53; P53; Cellular tumor antigen p53; Antigen NY-CO-13; Phosphoprotein p53; Tumor suppressor p53; anti-p53 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-p53 antibody	WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-p53 antibody	Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA- Mkl1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seems to have an effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-ARNTL/BMAL1- mediated transcriptional activation of PER2 (PubMed:24051492).
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	120407068
NCBI GeneID	7157
NCBI Accession #	NP_000537.3
NCBI GenBank Nucleotide #	NM_000546.5
UniProt Accession #	P04637; Q15086; Q15087; Q15088; Q16535; Q16807; Q16808; Q16809; Q16810; Q16811; Q16848; Q2XN98
Molecular Weight	43,653 Da
NCBI Official Full Name	cellular tumor antigen p53 isoform a
NCBI Official Synonym Full Names	tumor protein p53
NCBI Official Symbol	TP53
NCBI Official Synonym Symbols	P53; BCC7; LFS1; TRP53
NCBI Protein Information	cellular tumor antigen p53; antigen NY-CO-13; phosphoprotein p53; p53 tumor suppressor; transformation-related protein 53
UniProt Protein Name	Cellular tumor antigen p53
UniProt Gene Name	TP53
UniProt Synonym Gene Names	P53
UniProt Entry Name	P53_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/p53-antibody-western-blot-elisa/295541>