

Rabbit PCNA Polyclonal Antibody | anti-PCNA antibody

PCNA Polyclonal Antibody

CATALOG # AAA295563	HOST Rabbit	REACTIVITY Human, Mouse, Rat, Monkey	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PCNA; Polyclonal Antibody; PCNA Polyclonal Antibody; Proliferating cell nuclear antigen; Cyclin; anti-PCNA antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Applicable Applications for anti-PCNA antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-PCNA antibody

Auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic- apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage tolerance pathways (PubMed:24939902). Acts as a loading platform to recruit DDR proteins that allow completion of DNA replication after DNA damage and promote postreplication repair: Monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination mechanisms to synthesize across the lesion (PubMed:24695737).

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505641
NCBI GeneID	5111
NCBI Accession #	NP_002583.1
NCBI GenBank Nucleotide #	NM_002592.2
UniProt Accession #	P12004; B2R897; D3DW02
Molecular Weight	28,769 Da
NCBI Official Full Name	proliferating cell nuclear antigen
NCBI Official Synonym Full Names	proliferating cell nuclear antigen
NCBI Official Symbol	PCNA
NCBI Protein Information	proliferating cell nuclear antigen; cyclin; DNA polymerase delta auxiliary protein
UniProt Protein Name	Proliferating cell nuclear antigen
UniProt Gene Name	PCNA
UniProt Synonym Gene Names	PCNA
UniProt Entry Name	PCNA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/pcna-antibody-immunohistochemistry-western-blot-elisa/295563>