

Rabbit anti-Human CD32-B/C Polyclonal Antibody | anti-CD32-B/C antibody

CD32-B/C Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA296187	Rabbit	Human	Western Blot


PRODUCT SYNONYMS

CD32-B/C; Polyclonal Antibody; CD32-B/C Polyclonal Antibody; FCGR2B; CD32; FCG2; IGFR2; Low affinity immunoglobulin gamma Fc region receptor II-b; IgG Fc receptor II-b; CDw32; Fc-gamma RII-b; Fc-gamma-RIIb; FcRII-b; FCGR2C; IGFR2; Low affinity immunoglobulin gamma Fc region receptor II-c; IgG Fc receptor II-c; Fc-gamma RII-c; Fc-gamma-RIIc; FcRII-c; anti-CD32-B/C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD32-B/C antibody	WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CD32-B/C antibody

Receptor for the Fc region of complexed or aggregated immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B- cells. Binding to this receptor results in down-modulation of previous state of cell activation triggered via antigen receptors on B-cells (BCR), T-cells (TCR) or via another Fc receptor. Isoform IIB1 fails to mediate endocytosis or phagocytosis. Isoform IIB2 does not trigger phagocytosis.

NCBI AND UNIPROT INFORMATION

NCBI GI #	50511932
-----------	----------

NCBI GeneID	2213
NCBI Accession #	NP_001002273.1
NCBI GenBank Nucleotide #	NM_001002273.2
UniProt Accession #	P31994; O95649; Q53X85; Q5VXA9; Q8NIA1; A6H8N3
Molecular Weight	34,044 Da
NCBI Official Full Name	low affinity immunoglobulin gamma Fc region receptor II-b isoform 2
NCBI Official Synonym Full Names	Fc fragment of IgG, low affinity IIb, receptor (CD32)
NCBI Official Symbol	FCGR2B
NCBI Official Synonym Symbols	CD32; FCG2; CD32B; FCGR2; IGFR2
NCBI Protein Information	low affinity immunoglobulin gamma Fc region receptor II-b; CDw32; fcRII-b; Fc gamma RIIB; fc-gamma-RIIB; fc-gamma RII-b; igG Fc receptor II-b; Fc fragment of IgG, low affinity II, receptor for (CD32); Fc fragment of IgG, low affinity IIb, receptor for (CD
UniProt Protein Name	Low affinity immunoglobulin gamma Fc region receptor II-b
UniProt Gene Name	FCGR2B
UniProt Synonym Gene Names	CD32; FCG2; IGFR2; IgG Fc receptor II-b; Fc-gamma-RIIB; FcRII-b
UniProt Entry Name	FCG2B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd32-b-c-antibody-western-blot-elisa/296187>