

Rabbit MCM7 Polyclonal Antibody | anti-MCM7 antibody

MCM7 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA296242	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MCM7; Polyclonal Antibody; MCM7 Polyclonal Antibody; CDC47; MCM2; DNA replication licensing factor MCM7; CDC47 homolog; P1.1-MCM3; anti-MCM7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-MCM7 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MCM7 antibody	Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for S-phase checkpoint activation upon UV-induced damage.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	518830597
NCBI GeneID	4176
NCBI Accession #	NP_001265524.1
NCBI GenBank Nucleotide #	NM_001278595.1
UniProt Accession #	P33993; Q15076; Q96D34; Q96GL1; A4D2A1; A4D2A2; E9PGN9
Molecular Weight	719
NCBI Official Full Name	DNA replication licensing factor MCM7 isoform 2
NCBI Official Synonym Full Names	minichromosome maintenance complex component 7
NCBI Official Symbol	MCM7
NCBI Official Synonym Symbols	MCM2; CDC47; P85MCM; P1CDC47; PNAS146; P1.1-MCM3
NCBI Protein Information	DNA replication licensing factor MCM7; CDC47 homolog; homolog of S. cerevisiae Cdc47; minichromosome maintenance deficient 7
UniProt Protein Name	DNA replication licensing factor MCM7
UniProt Gene Name	MCM7
UniProt Synonym Gene Names	CDC47; MCM2
UniProt Entry Name	MCM7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mcm7-antibody-immunohistochemistry-western-blot-elisa/296242>