

Rabbit anti-Human CD107b Polyclonal Antibody | anti-CD107b antibody

CD107b Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA296436	Rabbit	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD107b; Polyclonal Antibody; CD107b Polyclonal Antibody; LAMP2; Lysosome-associated membrane glycoprotein 2; LAMP-2; Lysosome-associated membrane protein 2; CD107 antigen-like family member B; anti-CD107b antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD107b antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CD107b antibody	Plays an important role in chaperone-mediated autophagy, a process that mediates lysosomal degradation of proteins in response to various stresses and as part of the normal turnover of proteins with a long biological half-life (PubMed:11082038, PubMed:18644871, PubMed:24880125, PubMed:27628032). Functions by binding target proteins, such as GAPDH and MLLT11, and targeting them for lysosomal degradation (PubMed:11082038, PubMed:18644871, PubMed:24880125). Plays a role in lysosomal protein degradation in response to starvation (By similarity). Required for the fusion of autophagosomes with lysosomes during autophagy (PubMed:27628032). Cells that lack LAMP2 express normal levels of VAMP8, but fail to accumulate STX17 on autophagosomes, which is the most likely explanation for the lack of fusion between autophagosomes and lysosomes (PubMed:27628032). Required for normal degradation of the contents of autophagosomes (PubMed:27628032). Required for efficient MHCII-mediated presentation of exogenous antigens via its function in lysosomal protein degradation; antigenic peptides generated by proteases in the endosomal/lysosomal compartment are captured by nascent MHCII subunits (PubMed:20518820). Is not required for efficient MHCII-mediated presentation of endogenous antigens (PubMed:20518820).
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	1708854
NCBI GeneID	3920
UniProt Accession #	P13473; Q16641; Q6Q3G8; Q96J30; Q99534; Q9UD93; A8K4X5; D3DTF0
Molecular Weight	410
NCBI Official Full Name	Lysosome-associated membrane glycoprotein 2
NCBI Official Synonym Full Names	lysosomal-associated membrane protein 2
NCBI Official Symbol	LAMP2
NCBI Official Synonym Symbols	LAMPB; CD107b; LAMP-2; LGP110
NCBI Protein Information	lysosome-associated membrane glycoprotein 2; CD107 antigen-like family member B
UniProt Protein Name	Lysosome-associated membrane glycoprotein 2
UniProt Gene Name	LAMP2
UniProt Synonym Gene Names	LAMP-2; Lysosome-associated membrane protein 2
UniProt Entry Name	LAMP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd107b-antibody-immunohistochemistry-western-blot-elisa/296436>