

Rabbit MEK-1 Polyclonal Antibody | anti-MEK-1 antibody

MEK-1 Polyclonal Antibody

CATALOG # AAA296481	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

MEK-1; Polyclonal Antibody; MEK-1 Polyclonal Antibody; MAP2K1 MEK1 PRKMK1; anti-MEK-1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-MEK-1 antibody	WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: WB 1:500-2000, ELISA 1:10000-20000


WB (Western Blot)

Dilution: WB 1:500-2000, ELISA 1:10000-20000

ADDITIONAL INFORMATION

Related Product Information for anti-MEK-1 antibody	Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2. Both MAP2K1/MEK1 and MAP2K2/MEK2 function specifically in the MAPK/ERK cascade, and catalyze the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in the extracellular signal-regulated kinases MAPK3/ERK1 and MAPK1/ERK2, leading to their activation and further transduction of the signal within the MAPK/ERK cascade. Depending on the cellular context, this pathway mediates diverse biological functions such as cell growth, adhesion, survival and differentiation, predominantly through the regulation of transcription, metabolism and cytoskeletal rearrangements. One target of the MAPK/ERK cascade is peroxisome proliferator- activated receptor gamma (PPARG), a nuclear receptor that promotes differentiation and apoptosis. MAP2K1/MEK1 has been shown to export PPARG from the nucleus. The MAPK/ERK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	5579478
NCBI GeneID	5604
NCBI Accession #	NP_002746.1
NCBI GenBank Nucleotide #	NM_002755.3
UniProt Accession #	Q02750
Molecular Weight	43,439 Da
NCBI Official Full Name	dual specificity mitogen-activated protein kinase kinase 1
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 1
NCBI Official Symbol	MAP2K1
NCBI Official Synonym Symbols	MEK1; MKK1; MAPKK1; PRKMK1
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 1; MEK 1; MAPKK 1; MAPK/ERK kinase 1; ERK activator kinase 1; protein kinase, mitogen-activated, kinase 1 (MAP kinase kinase 1)
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 1
UniProt Gene Name	MAP2K1
UniProt Synonym Gene Names	MEK1; PRKMK1; MAP kinase kinase 1; MAPKK 1; MKK1; MEK 1
UniProt Entry Name	MP2K1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mek-1-antibody-western-blot-elisa/296481>