

Rabbit GRP78 BiP Polyclonal Antibody | anti-GRP78BiP antibody

GRP78 BiP Polyclonal Antibody

CATALOG # AAA296515	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

GRP78 BiP; Polyclonal Antibody; GRP78 BiP Polyclonal Antibody; 78 kDa glucose-regulated protein (GRP-78)(Endoplasmic reticulum luminal Ca (2+)-binding protein grp78)(Heat shock 70 kDa protein 5)(Immunoglobulin heavy chain-binding protein)(BiP); anti-GRP78BiP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-GRP78BiP antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: WB 1:500-2000, IHC-p 1:500-200, ELISA 1:10000-20000




IHC (Immunohistochemistry)

Dilution: WB 1:500-2000, IHC-p 1:500-200, ELISA 1:10000-20000



IHC (Immunohistochemistry)

Dilution: WB 1:500-2000, IHC-p 1:500-200, ELISA 1:10000-20000



WB (Western Blot)

Dilution: WB 1:500-2000, IHC-p 1:500-200, ELISA 1:10000-20000

ADDITIONAL INFORMATION

Related Product Information for anti-GRP78BiP antibody	Endoplasmic reticulum chaperone that plays a key role in protein folding and quality control in the endoplasmic reticulum lumen (PubMed:23769672, PubMed:23990668, PubMed:28332555). Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10/ERdj5, probably to facilitate the release of DNAJC10/ERdj5 from its substrate (By similarity). Acts as a key repressor of the ERN1/IRE1-mediated unfolded protein response (UPR)(PubMed:19538957). In the unstressed endoplasmic reticulum, recruited by DNAJB9/ERdj4 to the luminal region of ERN1/IRE1, leading to disrupt the dimerization of ERN1/IRE1, thereby inactivating ERN1/IRE1 (By similarity). Accumulation of misfolded protein in the endoplasmic reticulum causes release of HSPA5/BiP from ERN1/IRE1, allowing homodimerization and subsequent activation of ERN1/IRE1 (By similarity). Plays an auxiliary role in post-translational transport of small presecretory proteins across endoplasmic reticulum (ER). May function as an allosteric modulator for SEC61 channel-forming translocon complex, likely cooperating with SEC62 to enable the productive insertion of these precursors into SEC61 channel. Appears to specifically regulate translocation of precursors having inhibitory residues in their mature region that weaken channel gating.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	16507237
NCBI GeneID	3309
NCBI Accession #	NP_005338.1
NCBI GenBank Nucleotide #	NM_005347.4
UniProt Accession #	P11021; Q2EF78; Q9NPF1; Q9UK02; B0QZ61
Molecular Weight	72,333 Da
NCBI Official Full Name	78 kDa glucose-regulated protein
NCBI Official Synonym Full Names	heat shock 70kDa protein 5 (glucose-regulated protein, 78kDa)
NCBI Official Symbol	HSPA5
NCBI Official Synonym Symbols	BIP; MIF2; GRP78
NCBI Protein Information	78 kDa glucose-regulated protein; immunoglobulin heavy chain-binding protein; endoplasmic reticulum luminal Ca(2+)-binding protein grp78
UniProt Protein Name	78 kDa glucose-regulated protein
UniProt Gene Name	HSPA5
UniProt Synonym Gene Names	GRP78; GRP-78; BiP
UniProt Entry Name	GRP78_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/grp78-bip-antibody-immunohistochemistry-western-blot-elisa/296515>