

Rabbit anti-Human CD3 zeta Polyclonal Antibody | anti-CD3zeta antibody

CD3 zeta Polyclonal Antibody

CATALOG # AAA296631	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CD3 zeta; Polyclonal Antibody; CD3 zeta Polyclonal Antibody; T-cell surface glycoprotein CD3 zeta chain (T-cell receptor T3 zeta chain)(CD antigen CD247); anti-CD3zeta antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD3zeta antibody	IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES



IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000

ADDITIONAL INFORMATION

Related Product Information for anti-CD3zeta antibody	Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:7509083). CD3Z ITAMs phosphorylation creates multiple docking sites for the protein kinase ZAP70 leading to ZAP70 phosphorylation and its conversion into a catalytically active enzyme (PubMed:7509083). Plays an important role in intrathymic T-cell differentiation. Additionally, participates in the activity-dependent synapse formation of retinal ganglion cells (RGCs) in both the retina and dorsal lateral geniculate nucleus (dLGN)(By similarity).
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557431
NCBI GeneID	919
NCBI Accession #	NP_000725.1
NCBI GenBank Nucleotide #	NM_000734.3
UniProt Accession #	P20963; Q5VX13; Q8TAX4; B1AK49; A46461
Molecular Weight	18,696 Da
NCBI Official Full Name	T-cell surface glycoprotein CD3 zeta chain isoform 2
NCBI Official Synonym Full Names	CD247 molecule
NCBI Official Symbol	CD247
NCBI Official Synonym Symbols	T3Z; CD3H; CD3Q; CD3Z; TCRZ; CD3-ZETA
NCBI Protein Information	T-cell surface glycoprotein CD3 zeta chain; CD3zeta chain; TCR zeta chain; CD247 antigen, zeta subunit; T-cell receptor T3 zeta chain; CD3Z antigen, zeta polypeptide (TiT3 complex); T-cell antigen receptor complex, zeta subunit of CD3
UniProt Protein Name	T-cell surface glycoprotein CD3 zeta chain
UniProt Gene Name	CD247
UniProt Synonym Gene Names	CD3Z; T3Z; TCRZ
UniProt Entry Name	CD3Z_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.