

Rabbit Frizzled-4 Polyclonal Antibody | anti-FZD4 antibody

Frizzled-4 Polyclonal Antibody

CATALOG # AAA296639	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Frizzled-4; Polyclonal Antibody; Frizzled-4 Polyclonal Antibody; Frizzled-4 (Fz-4)(hFz4)(FzE4)(CD antigen CD344); anti-FZD4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-FZD4 antibody	IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES



IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000

ADDITIONAL INFORMATION

Related Product Information for anti-FZD4 antibody	Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin (CTNNB1) canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin (CTNNB1) and activation of Wnt target genes. Plays a critical role in retinal vascularization by acting as a receptor for Wnt proteins and norrin (NDP). In retina, it can be both activated by Wnt protein-binding, but also by a Wnt-independent signaling via binding of norrin (NDP), promoting in both cases beta-catenin (CTNNB1) accumulation and stimulation of LEF/TCF-mediated transcriptional programs. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	22547161
NCBI GeneID	8322
NCBI Accession #	NP_036325.2
NCBI GenBank Nucleotide #	NM_012193.3
UniProt Accession #	Q9ULV1; Q14C97; Q6S9E4; A8K9Q3
Molecular Weight	59,881 Da
NCBI Official Full Name	frizzled-4
NCBI Official Synonym Full Names	frizzled family receptor 4
NCBI Official Symbol	FZD4
NCBI Official Synonym Symbols	Fz4; EVR1; FEVR; Fz-4; FzE4; GPCR; hFz4; CD344; FZD4S
NCBI Protein Information	frizzled-4; frizzled homolog 4; WNT receptor frizzled-4; frizzled 4, seven transmembrane spanning receptor
UniProt Protein Name	Frizzled-4
UniProt Gene Name	FZD4
UniProt Synonym Gene Names	Fz-4; hFz4
UniProt Entry Name	FZD4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/frizzled-4-antibody-immunohistochemistry-elisa/296639>