

Rabbit C/EBP beta Polyclonal Antibody | anti-C/EBPbeta antibody

C/EBP beta Polyclonal Antibody

CATALOG # AAA296681	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

C/EBP beta; Polyclonal Antibody; C/EBP beta Polyclonal Antibody; CEBPB; LAP; TCF5; PP9092; CCAAT/enhancer-binding protein beta; Liver activator protein; Nuclear factor NF-IL6; Transcription factor 5; TCF-5; anti-C/EBPbeta antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-C/EBPbeta antibody	WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: WB 1:500-2000, ELISA 1:10000-20000



WB (Western Blot)

Dilution: WB 1:500-2000, ELISA 1:10000-20000

ADDITIONAL INFORMATION

Related Product Information for anti-C/EBPbeta antibody	Important transcription factor regulating the expression of genes involved in immune and inflammatory responses (PubMed:9374525, PubMed:12048245, PubMed:18647749). Plays also a significant role in adipogenesis, as well as in the gluconeogenic pathway, liver regeneration, and hematopoiesis. The consensus recognition site is 5'-T[TG]NNGNAA[TG]-3'. Its functional capacity is governed by protein interactions and post-translational protein modifications. During early embryogenesis, plays essential and redundant functions with CEBPA. Has a promotitic effect on many cell types such as hepatocytes and adipocytes but has an antiproliferative effect on T-cells by repressing MYC expression, facilitating differentiation along the T-helper 2 lineage. Binds to regulatory regions of several acute-phase and cytokines genes and plays a role in the regulation of acute-phase reaction and inflammation. Plays also a role in intracellular bacteria killing (By similarity). During adipogenesis, is rapidly expressed and, after activation by phosphorylation, induces CEBPA and PPARG, which turn on the series of adipocyte genes that give rise to the adipocyte phenotype. The delayed transactivation of the CEBPA and PPARG genes by CEBPB appears necessary to allow mitotic clonal expansion and thereby progression of terminal differentiation (PubMed:20829347). Essential for female reproduction because of a critical role in ovarian follicle development (By similarity). Restricts osteoclastogenesis: together with NFE2L1; represses expression of DSPP during odontoblast differentiation (By similarity).
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	28872796
NCBI GeneID	1051
NCBI Accession #	NP_005185.2
NCBI GenBank Nucleotide #	NM_005194.3
UniProt Accession #	P17676; Q96IH2; Q9H4Z5; A8K671
Molecular Weight	36,106 Da
NCBI Official Full Name	CCAAT/enhancer-binding protein beta
NCBI Official Synonym Full Names	CCAAT/enhancer binding protein (C/EBP), beta
NCBI Official Symbol	CEBPB
NCBI Official Synonym Symbols	LAP; CRP2; TCF5; IL6DBP; NF-IL6; C/EBP-beta
NCBI Protein Information	CCAAT/enhancer-binding protein beta; TCF-5; nuclear factor NF-IL6; transcription factor 5; nuclear factor of interleukin 6; interleukin 6-dependent DNA-binding protein; liver-enriched transcriptional activator protein
UniProt Protein Name	CCAAT/enhancer-binding protein beta
UniProt Gene Name	CEBPB
UniProt Synonym Gene Names	LAP; TCF5; C/EBP beta; TCF-5
UniProt Entry Name	CEBPB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.