

Rabbit N Cadherin Monoclonal Antibody | anti-NCAD antibody

N Cadherin Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA30054	Rabbit	Human, Mouse, Rat	Western Blot, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

N Cadherin; Monoclonal Antibody; N Cadherin Antibody; CADH2_HUMAN antibody; Cadherin 2 antibody; Cadherin 2 N cadherin neuronal antibody; Cadherin 2 type 1 antibody; Cadherin 2 type 1 N cadherin neuronal antibody; Cadherin 2; type 1; N-cadherin (neuronal) antibody; Cadherin-2 antibody; Cadherin2 antibody; Calcium dependent adhesion protein neuronal antibody; CD325 antibody; CD325 antigen antibody; CDH2 antibody; CDHN antibody; CDw325 antibody; CDw325 antigen antibody; N cadherin 1 antibody; N-cadherin antibody; NCAD antibody; Neural cadherin antibody; OTTHUMP00000066304 antibody; OTTHUMP00000067378 antibody; anti-NCAD antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Clone Number	SY02-46
Purity/Purification	ProA affinity purified
Form/Format	1*TBS (pH7.4), 1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Applicable Applications for anti-NCAD antibody	WB (Western Blot), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry)
Application Notes	WB: 1:1000-5000 IHC: 1:50-1:200 ICC: 1:50-1:200 FC/FACS: 1:50-1:100
Antibody Type	Recombinant Antibody
Immunogen	Recombinant protein
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C.




FCM (Flow Cytometry)

Flow cytometric analysis of Hela cells with N Cadherin antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.




ICC (Immunocytochemistry)

ICC staining N Cadherin in RH-35 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




ICC (Immunocytochemistry)

ICC staining N Cadherin in NCCIT cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




ICC (Immunocytochemistry)

ICC staining N Cadherin in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC (Immunocytochemistry)

ICC staining N Cadherin in F9 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded mouse heart tissue using anti-N Cadherin antibody. Counter stained with hematoxylin.

ADDITIONAL INFORMATION

Related Product Information for anti-NCAD antibody

Cadherins comprise a family of Ca²⁺-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. The classical cadherins, E-, N- and P-cadherin, consist of large extracellular domains characterized by a series of five homologous NH₂ terminal repeats. The most distal of these cadherins is thought to be responsible for binding specificity, transmembrane domains and carboxy-terminal intracellular domains. The relatively short intracellular domains interact with a variety of cytoplasmic proteins, such as b-catenin, to regulate cadherin function. Members of this family of adhesion proteins include rat cadherin K (and its human homolog, cadherin-6), R-cadherin, B-cadherin, E/P cadherin and cadherin-5.

Product Categories/Family for anti-NCAD antibody	Total protein Ab
--	------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	14589889
NCBI GeneID	1000
NCBI Accession #	NP_001783.2
NCBI GenBank Nucleotide #	NM_001792.3
UniProt Accession #	P19022; Q14923; Q8N173; B0YIY6
Molecular Weight	99,809 Da
NCBI Official Full Name	cadherin-2 preproprotein
NCBI Official Synonym Full Names	cadherin 2, type 1, N-cadherin (neuronal)
NCBI Official Symbol	CDH2
NCBI Official Synonym Symbols	CDHN; NCAD; CD325; CDw325
NCBI Protein Information	cadherin-2; N-cadherin 1; neural cadherin; neural-cadherin; cadherin 2, N-cadherin (neuronal); calcium-dependent adhesion protein, neuronal
UniProt Protein Name	Cadherin-2
UniProt Gene Name	CDH2
UniProt Synonym Gene Names	CDHN; NCAD; N-cadherin
UniProt Entry Name	CADH2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/n-cadherin-antibody-clone-sy02-46-western-blot-immunocytochemistry-immunofluorescence-immunohistochemistry-flow-cytometry-elisa/30054>