

Rabbit PSMA1 Polyclonal Antibody | anti-PSMA1 antibody

PSMA1 Antibody

CATALOG # AAA301999	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PSMA1; Polyclonal Antibody; PSMA1 Antibody; NU; HC2; PROS30; anti-PSMA1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total PSMA1 protein.
Purity/Purification	Antibodies were purified by affinity purification using immunogen.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	263
Applicable Applications for anti-PSMA1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Recombinant Protein
Immunogen Description	Recombinant protein of human PSMA1.
Target Name	PSMA1
Preparation and Storage	Store at -20 degree C


IMAGES


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded rat lung using PSMA1 antibody at dilution of 1:100 (200x lens).



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human stomach cancer using PSMA1 antibody at dilution of 1:100 (400x lens).




IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded mouse heart using PSMA1 antibody at dilution of 1:100 (400x lens).



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast cancer using PSMA1 antibody at dilution of 1:100 (400x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using PSMA1 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-PSMA1 antibody

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Product Categories/Family for anti-PSMA1 antibody

Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506179
NCBI GeneID	5682
NCBI Accession #	NP_002777.1
NCBI GenBank Nucleotide #	NM_002786.3
UniProt Accession #	P25786; Q53YE8; Q9BRV9; A8K400
Molecular Weight	30,239 Da
NCBI Official Full Name	proteasome subunit alpha type-1 isoform 2
NCBI Official Synonym Full Names	proteasome subunit alpha 1
NCBI Official Symbol	PSMA1
NCBI Official Synonym Symbols	NU; HC2; PROS30; HEL-S-275

NCBI Protein Information	proteasome subunit alpha type-1
UniProt Protein Name	Proteasome subunit alpha type-1
UniProt Gene Name	PSMA1
UniProt Synonym Gene Names	HC2; NU; PROS30; PSC2; PROS-30
UniProt Entry Name	PSA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/psma1-antibody-immunohistochemistry-western-blot-elisa/301999>