

Rabbit anti-Human STAT1 Polyclonal Antibody | anti-STAT1 antibody

STAT1 (Ab-701) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA302529	Rabbit	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

STAT1; Polyclonal Antibody; STAT1 (Ab-701) Antibody; CANDF7; ISGF-3; STAT91; anti-STAT1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total STAT1 protein.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	750
Applicable Applications for anti-STAT1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa. 699~703 (T-G-Y-I-K) derived from Human STAT1.
Target Name	STAT1
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using STAT1(Ab-701) Antibody (left) or the same antibody preincubated with blocking peptide(right).



WB (Western Blot)

Western blot analysis of extracts from HeLa, A431 and JK cells using STAT1(Ab-701) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-STAT1 antibody	Signal transducer and activator of transcription that mediates signaling by interferons (IFNs). Following type I IFN (IFN-a and IFN-beta) binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state. In response to type II IFN (IFN-gamma), STAT1 is tyrosine- and serine-phosphorylated. It then forms a homodimer termed IFN-gamma-activated factor (GAF), migrates into the nucleus and binds to the IFN gamma activated sequence (GAS) to drive the expression of the target genes, inducing a cellular antiviral state.
Product Categories/Family for anti-STAT1 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	6274552
NCBI GeneID	6772
NCBI Accession #	NP_009330.1
NCBI GenBank Nucleotide #	NM_007315.3
UniProt Accession #	P42224; Q53S88; Q53XW4; Q68D00; Q9UDL5; A8K989; B2RCA0; D2KFR8; D3DPI7
Molecular Weight	83,043 Da
NCBI Official Full Name	signal transducer and activator of transcription 1-alpha/beta isoform alpha
NCBI Official Synonym Full Names	signal transducer and activator of transcription 1, 91kDa
NCBI Official Symbol	STAT1
NCBI Official Synonym Symbols	CANDF7; IMD31A; IMD31B; IMD31C; ISGF-3; STAT91
NCBI Protein Information	signal transducer and activator of transcription 1-alpha/beta
UniProt Protein Name	Signal transducer and activator of transcription 1-alpha/beta
UniProt Gene Name	STAT1
UniProt Entry Name	STAT1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/stat1-antibody-immunohistochemistry-western-blot-elisa/302529>