

Rabbit anti-Human c-Cbl Polyclonal Antibody | anti-c-Cbl antibody

c-Cbl (Ab-700) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA302545	Rabbit	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

c-Cbl; Polyclonal Antibody; c-Cbl (Ab-700) Antibody; Signal transduction protein CBL; Proto-oncogene c-CBL; Casitas B-lineage lymphoma proto-oncogene; RING finger protein 55; anti-c-Cbl antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total c-Cbl protein.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	906
Applicable Applications for anti-c-Cbl antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa. 698-702 (T-E-Y-M-T) derived from Human c-Cbl.
Target Name	c-Cbl
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using c-Cbl(Ab-700) Antibody (left) or the same antibody preincubated with blocking peptide(right).



WB (Western Blot)

Western blot analysis of extracts from HeLa and HT29 cells using c-Cbl(Ab-700) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-c-Cbl antibody	Participates in signal transduction in hematopoietic cells. Adapter protein that functions as a negative regulator of many signaling pathways that start from receptors at the cell surface. Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including PDGFA, EGF and CSF1, and terminates signaling.
Product Categories/Family for anti-c-Cbl antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	52426745
NCBI GeneID	867
NCBI Accession #	NP_005179.2
NCBI GenBank Nucleotide #	NM_005188.3
UniProt Accession #	P22681; A3KMP8
Molecular Weight	99,633 Da
NCBI Official Full Name	E3 ubiquitin-protein ligase CBL
NCBI Official Synonym Full Names	Cbl proto-oncogene, E3 ubiquitin protein ligase
NCBI Official Symbol	CBL
NCBI Official Synonym Symbols	CBL2; NSLL; C-CBL; RNF55; FRA11B
NCBI Protein Information	E3 ubiquitin-protein ligase CBL
UniProt Protein Name	E3 ubiquitin-protein ligase CBL
UniProt Gene Name	CBL
UniProt Synonym Gene Names	CBL2; RNF55
UniProt Entry Name	CBL_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/c-cbl-antibody-immunohistochemistry-western-blot-elisa/302545>