

Rabbit anti-Human, Mouse ADAM15 Polyclonal Antibody | anti-ADAM15 antibody

ADAM15 Antibody

CATALOG # AAA302762	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	-----------------------------------	---


PRODUCT SYNONYMS

ADAM15; Polyclonal Antibody; ADAM15 Antibody; MDC15; anti-ADAM15 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of total ADAM15 protein.
Purity/Purification	Antigen affinity purification.
Form/Format	Rabbit IgG in pH7.3 PBS, 0.05% NaN ₃ , 50% Glycerol.
Concentration	1.5 mg/ml (varies by lot)
Sequence Length	824
Applicable Applications for anti-ADAM15 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Recombinant Protein
Immunogen Description	Fusion protein corresponding to a region derived from internal residues of human ADAM metallopeptidase domain 15
Target Name	ADAM15
Preparation and Storage	Store at -20 degree C

IMAGES


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human lung cancer tissue using at dilution 1/40.


Application Data

Gel: 8%SDS-PAGE Lysates (from left to right): Human liver cancer tissue, NIH/3T3 and Jurkat cell Amount of lysate: 40ug per lane Primary antibody: 1/300 dilution Secondary antibody dilution: 1/8000 Exposure time: 5 minutes

ADDITIONAL INFORMATION

Related Product Information for anti-ADAM15 antibody	The protein encoded by this gene is a member of the ADAM (a disintegrin and metalloproteinase) protein family. ADAM family members are type I transmembrane glycoproteins known to be involved in cell adhesion and proteolytic ectodomain processing of cytokines and adhesion molecules. This protein contains multiple functional domains including a zinc-binding metalloprotease domain, a disintegrin-like domain, as well as a EGF-like domain. Through its disintegrin-like domain, this protein specifically interacts with the integrin beta chain, beta 3. It also interacts with Src family protein-tyrosine kinases in a phosphorylation-dependent manner, suggesting that this protein may function in cell-cell adhesion as well as in cellular signaling. Multiple alternatively spliced transcript variants encoding distinct isoforms have been observed.
Product Categories/Family for anti-ADAM15 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	387849014
NCBI GeneID	8751
NCBI Accession #	NP_001248393.1
NCBI GenBank Nucleotide #	NM_001261464.1
UniProt Accession #	Q13444; Q13493; Q53XQ0; Q5SR68; Q5SR69; Q6R267; Q71S61; Q71S62; B3KQU5; B4DLB5; B4DMH8; E9PN65
Molecular Weight	68,256 Da
NCBI Official Full Name	disintegrin and metalloproteinase domain-containing protein 15 isoform 7 preproprotein
NCBI Official Synonym Full Names	ADAM metalloproteinase domain 15
NCBI Official Symbol	ADAM15
NCBI Official Synonym Symbols	MDC15
NCBI Protein Information	disintegrin and metalloproteinase domain-containing protein 15
UniProt Protein Name	Disintegrin and metalloproteinase domain-containing protein 15
UniProt Gene Name	ADAM15
UniProt Synonym Gene Names	MDC15; ADAM 15; MDC-15
UniProt Entry Name	ADA15_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/adam15-antibody-immunohistochemistry-western-blot-elisa/302762>