

Rabbit anti-Human C1QBP Polyclonal Antibody | anti-C1QBP antibody

C1QBP Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA302894	Rabbit	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

C1QBP; Polyclonal Antibody; C1QBP Antibody; C1q globular domain-binding protein; complement component 1; q subcomponent binding protein; Complement component 1; Q subcomponent binding protein; anti-C1QBP antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of total C1QBP protein.
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	282
Applicable Applications for anti-C1QBP antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide
Immunogen Description	Synthesized peptide derived from internal of human C1QBP.
Target Name	C1QBP
Preparation and Storage	Store at -20 degree C

IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry analysis of paraffin-embedded human tonsil tissue, using C1QBP antibody.



WB (Western Blot)
Western blot analysis of extracts from Jurkat cells, K562 cells and HUVEC cells, using C1QBP antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-C1QBP antibody	Is believed to be a multifunctional and multicompartmental protein involved in inflammation and infection processes, ribosome biogenesis, regulation of apoptosis, transcriptional regulation and pre-mRNA splicing. At the cell surface is thought to act as an endothelial receptor for plasma proteins of the complement and kallikrein-kinin cascades. Putative receptor for C1q; specifically binds to the globular "heads" of C1q thus inhibiting C1; may perform the receptor function through a complex with C1qR/CD93. In complex with cytokeratin-1/KRT1 is a high affinity receptor for kininogen-1/HMWK. Can also bind other plasma proteins, such as coagulation factor XII leading to its autoactivation. May function to bind initially fluid kininogen-1 to the cell membrane. The secreted form may enhance both extrinsic and intrinsic coagulation pathways. It is postulated that the cell surface form requires docking with transmembrane proteins for downstream signaling which might be specific for a cell-type or response. By acting as C1q receptor is involved in chemotaxis of immature dendritic cells and neutrophils and is proposed to signal through CD209/DC-SIGN on immature dendritic cells, through integrin alpha-4/beta-1 during trophoblast invasion of the decidua, and through integrin beta-1 during endothelial cell adhesion and spreading. Signaling involved in inhibition of innate immune response is implicating the PI3K-AKT/PKB pathway. In mitochondrial translation may be involved in formation of functional 55S mitoribosomes; the function seems to involve its RNA-binding activity. May be involved in the nucleolar ribosome maturation process; the function may involve the exchange of FBL for RRP1 in the association with pre-ribosome particles. Involved in regulation of RNA splicing by inhibiting the RNA-binding capacity of SRSF1 and its phosphorylation. Is required for the nuclear translocation of splicing factor U2AF1L4. Involved in regulation of CDKN2A- and HRK-mediated apoptosis. Stabilizes mitochondrial CDKN2A isoform smARF. May be involved in regulation of FOXC1 transcriptional activity and NFY/CCAAT-binding factor complex-mediated transcription. In infection processes acts as an attachment site for microbial proteins, including <i>Listeria monocytogenes</i> internalin B and <i>Staphylococcus aureus</i> protein A. May play a role in antibacterial defense as it can bind to cell surface hyaluronan and inhibit <i>Streptococcus pneumoniae</i> hyaluronate lyase. Involved in replication of Rubella virus. May be involved in modulation of the immune response; ligation by HCV core protein is resulting in suppression of interleukin-12 production in monocyte-derived dendritic cells. Involved in regulation of antiviral response by inhibiting DDX58- and IFIH1-mediated signaling pathways probably involving its association with MAVS after viral infection. Involved in HIV-1 replication, presumably by contributing to splicing of viral RNA. Honore B., Gene 134:283-287(1993). Ghebrehwet B., J. Exp. Med. 179:1809-1821(1994). Tye A.J., J. Biol. Chem. 276:17069-17075(2001)
Product Categories/Family for anti-C1QBP antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502491
NCBI GeneID	708
NCBI Accession #	NP_001203.1
NCBI GenBank Nucleotide #	NM_001212.3

UniProt Accession #	Q07021 ; Q2HXR8 ; Q9NNY8
Molecular Weight	31,362 Da
NCBI Official Full Name	complement component 1 Q subcomponent-binding protein, mitochondrial
NCBI Official Synonym Full Names	complement component 1, q subcomponent binding protein
NCBI Official Symbol	C1QBP
NCBI Official Synonym Symbols	p32; HABP1; gC1qR; GC1QBP; SF2p32; gC1Q-R
NCBI Protein Information	complement component 1 Q subcomponent-binding protein, mitochondrial
UniProt Protein Name	Complement component 1 Q subcomponent-binding protein, mitochondrial
UniProt Gene Name	C1QBP
UniProt Synonym Gene Names	GC1QBP; HABP1; SF2P32; C1qBP
UniProt Entry Name	C1QBP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/c1qbp-antibody-immunohistochemistry-western-blot-elisa/302894>