

Rabbit anti-Human Androgen Receptor Polyclonal Antibody | anti-AR antibody

Androgen Receptor (Phospho-Ser213) Antibody

CATALOG # AAA302949	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

Androgen Receptor; Polyclonal Antibody; Androgen Receptor (Phospho-Ser213) Antibody; ANDR; DHTR; AR; anti-AR antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	920
Applicable Applications for anti-AR antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 213 (E-A-S(p)-G-A) derived from Human Androgen Receptor.
Modification	Phospho-Ser213
Target Name	Androgen Receptor
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES



IF (Immunofluorescence)
Immunofluorescence staining of methanol-fixed HeLa cells using Androgen Receptor(Phospho-Ser213) Antibody.



WB (Western Blot)
Western blot analysis of extracts from DU145 cells using Androgen Receptor(Phospho-Ser213) Antibody (Lane 2) and the same antibody preincubated with blocking peptide (Lane 1).

ADDITIONAL INFORMATION

Related Product Information for anti-AR antibody	The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract causes spinal bulbar muscular atrophy (Kennedy disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Two alternatively spliced variants encoding distinct isoforms have been described.
Product Categories/Family for anti-AR antibody	Phospho-specific Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	21322252
NCBI GeneID	367
NCBI Accession #	NP_000035.2
NCBI GenBank Nucleotide #	NM_000044.3
UniProt Accession #	P10275; Q9UD95; A2RUN2; B1AKD7
Molecular Weight	44,643 Da
NCBI Official Full Name	androgen receptor isoform 1
NCBI Official Synonym Full Names	androgen receptor
NCBI Official Symbol	AR
NCBI Official Synonym Symbols	KD; AIS; AR8; TFM; DHTR; SBMA; HYSYP1; NR3C4; SMAX1; HUMARA
NCBI Protein Information	androgen receptor
UniProt Protein Name	Androgen receptor
UniProt Gene Name	AR
UniProt Synonym Gene Names	DHTR; NR3C4
UniProt Entry Name	ANDR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/androgen-receptor-antibody-immunofluorescence-western-blot-elisa/302949>