

Rabbit anti-Human XRCC5 Polyclonal Antibody | anti-XRCC5 antibody

XRCC5 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA302986	Rabbit	Human	Immunohistochemistry, Western Blot

PRODUCT SYNONYMS


XRCC5; Polyclonal Antibody; XRCC5 Antibody; 86 kDa subunit of Ku antigen; ATP-dependent DNA helicase II; 80 kDa subunit; CTC box binding factor 85 kDa subunit; CTC85; anti-XRCC5 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of total XRCC5 protein.
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	732
Applicable Applications for anti-XRCC5 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide
Immunogen Description	Synthesized peptide derived from C-terminal of human XRCC5.
Target Name	XRCC5
Preparation and Storage	Store at -20 degree C



IHC (Immunohistochemistry)
Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue using XRCC5 antibody.



WB (Western Blot)
Western blot analysis of extracts from Jurkat cells, using XRCC5 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-XRCC5 antibody	Single-stranded DNA-dependent ATP-dependent helicase. Has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by XRCC6. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The XRCC5/6 dimer acts as regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold. The XRCC5/6 dimer is probably involved in stabilizing broken DNA ends and bringing them together. The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step. In association with NAA15, the XRCC5/6 dimer binds to the osteocalcin promoter and activates osteocalcin expression. The XRCC5/6 dimer probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks. XRCC5 probably acts as the catalytic subunit of 5'-dRP activity, and allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined. The XRCC5/6 dimer together with APEX1 acts as a negative regulator of transcription.Ming-Ni Lee, Clin. Cancer Res., Feb 2007; 13: 832 - 838. PA Jeggo, PNAS, Jul 1992; 89: 6423. Patrick Hayden, Blood (ASH Annual Meeting Abstracts), Nov 2006; 108: 3416.
Product Categories/Family for anti-XRCC5 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	10863945
NCBI GeneID	7520
NCBI Accession #	NP_066964.1
NCBI GenBank Nucleotide #	NM_021141.3
UniProt Accession #	P13010; Q0Z7V0; Q4VBQ5; Q53HH7; Q7M4N0; Q9UCQ0; Q9UCQ1; A8K3X5
Molecular Weight	82,705 Da
NCBI Official Full Name	X-ray repair cross-complementing protein 5
NCBI Official Synonym Full Names	X-ray repair complementing defective repair in Chinese hamster cells 5 (double-strand-break rejoining)
NCBI Official Symbol	XRCC5
NCBI Official Synonym Symbols	KU80; KUB2; Ku86; NFIV; KARP1; KARP-1
NCBI Protein Information	X-ray repair cross-complementing protein 5

UniProt Protein Name	X-ray repair cross-complementing protein 5
UniProt Gene Name	XRCC5
UniProt Synonym Gene Names	G22P2; CTC85; CTCBF; TLAA
UniProt Entry Name	XRCC5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/xrcc5-antibody-immunohistochemistry-western-blot-elisa/302986>