

Rabbit ATF2 Polyclonal Antibody | anti-ATF2 antibody

ATF2 (Phospho-Thr71 or 53) Antibody

CATALOG # AAA303052	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

ATF2; Polyclonal Antibody; ATF2 (Phospho-Thr71 or 53) Antibody; CREB2; CREBP1; anti-ATF2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of ATF-2 only when phosphorylated at threonine 71 or 53.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	505
Applicable Applications for anti-ATF2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of threonine 71 or 53 (T-P-T(p)-P-T) derived from Human ATF2.
Modification	Phospho-Thr71 or 53
Target Name	ATF2
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using ATF2(Phospho-Thr71 or 53) Antibody (left) or the same antibody preincubated with blocking peptide(right).



WB (Western Blot)
Western blot analysis of extracts from HUVEC cells untreated or treated with Anisomycin using ATF2 (Phospho-Thr71 or 53) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-ATF2 antibody	Transcriptional activator, probably constitutive, which binds to the cAMP-responsive element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Interaction with JUN redirects JUN to bind to CRES preferentially over the 12-O-tetradecanoylphorbol-13-acetate response elements (TRES) as part of an ATF2-c-Jun complex.
Product Categories/Family for anti-ATF2 antibody	Phospho-specific Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	368711269
NCBI GeneID	1386
NCBI Accession #	NP_001243019.1
NCBI GenBank Nucleotide #	NM_001256090.1
UniProt Accession #	P15336; Q13000; Q3B7B7; Q4ZFU9; Q53RY2; Q8TAR1; A1L3Z2; A4D7U4; A4D7U5; A4D7V1; D3DPE9; G8JLM5
Molecular Weight	15,409 Da
NCBI Official Full Name	cyclic AMP-dependent transcription factor ATF-2 isoform 1
NCBI Official Synonym Full Names	activating transcription factor 2
NCBI Official Symbol	ATF2
NCBI Official Synonym Symbols	HB16; CREB2; TREB7; CREB-2; CRE-BP1
NCBI Protein Information	cyclic AMP-dependent transcription factor ATF-2
UniProt Protein Name	Cyclic AMP-dependent transcription factor ATF-2
UniProt Gene Name	ATF2
UniProt Synonym Gene Names	CREB2; CREBP1; cAMP-dependent transcription factor ATF-2; CREB-2; cAMP-responsive element-binding protein 2
UniProt Entry Name	ATF2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.