

Rabbit anti-Human, Mouse S100 Monoclonal Antibody | anti-S100 antibody

S100 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA30331	Rabbit	Human, Mouse	Western Blot, Immunoprecipitation, Immunohistochemistry, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS


S100; Monoclonal Antibody; S100 Antibody; NEF antibody; Protein S100 B antibody; Protein S100-B antibody; S 100 calcium binding protein beta chain antibody; S 100 protein beta chain antibody; S-100 protein beta chain antibody; S-100 protein subunit beta antibody; S100 antibody; S100 calcium binding protein beta (neural) antibody; S100 calcium-binding protein B antibody; S100 protein beta chain antibody; S100B antibody; S100B_HUMAN antibody; S100beta antibody; anti-S100 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Monoclonal
Clone Number	JM51-31
Purity/Purification	ProA affinity purified
Form/Format	1*TBS (pH7.4), 1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Applicable Applications for anti-S100 antibody	WB (Western Blot), IP (Immunoprecipitation), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry)
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200 IP: 1:10-1:50 FC/FACS: 1:50-1:100
Antibody Type	Recombinant Antibody
Immunogen	Recombinant protein
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C.




FCM (Flow Cytometry)
Flow cytometric analysis of SH-SY5Y cells with S100 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.




IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded mouse spinal cord tissue using anti-S100 antibody. Counter stained with hematoxylin.




IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti- S100 antibody. Counter stained with hematoxylin.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti- S100 antibody. Counter stained with hematoxylin.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti- S100 antibody. Counter stained with hematoxylin.



WB (Western Blot)
Western blot analysis of S100 on mouse heart cells lysates using anti-S100 antibody at 1/500 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-S100 antibody	The family of EF-hand type Ca2+-binding proteins includes calbindin (previously designated vitamin D-dependent Ca2+-binding protein), S-100 alpha and beta , calgranulins A (also designated MRP8), B (also designated MRP14) and C (S-100 like proteins), and the parvalbumin family members, including parvalbumin alpha and parvalbumin beta (also designated oncomodulin). The S-100 protein is involved in the regulation of cellular processes such as cell cycle progression and differentiation. Research also indicates that the S-100 protein may function in the activation of Ca2+ induced Ca2+ release, inhibition of microtubule assembly and inhibition of protein kinase C mediated phosphorylation. Two S-100 subunits, sharing 60% sequence identity, have been described as S-100 alpha chain and S-100 beta chain. Three S-100 dimeric forms have been characterized, differing in their subunit composition of either two alpha chains, two beta chains or one alpha and one beta chain. S-100 localizes to the cytoplasm and nuclei of astrocytes, Schwann's cells, ependymomas and astroglomas. S-100 is also detected in almost all benign naevi, malignant melanocytic tumours and in Langerhans cells in the skin. Calbindin, S-100 proteins and parvalbumin proteins are each expressed in neural tissues. In addition, S-100 alpha and beta are present in a variety of other tissues, and calbindin is present in intestine and kidney.
Product Categories/Family for anti-S100 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	5454032
NCBI GeneID	6271
NCBI Accession #	NP_006262.1
NCBI GenBank Nucleotide #	NM_006271.1
UniProt Accession #	P23297; Q5T7Y3; B2R5D9
Molecular Weight	10,546 Da
NCBI Official Full Name	protein S100-A1
NCBI Official Synonym Full Names	S100 calcium binding protein A1
NCBI Official Symbol	S100A1
NCBI Official Synonym Symbols	S100; S100A; S100-alpha
NCBI Protein Information	protein S100-A1; S100 alpha; S-100 protein alpha chain; S-100 protein subunit alpha; S100 calcium-binding protein A1; S100 protein, alpha polypeptide
UniProt Protein Name	Protein S100-A1
UniProt Gene Name	S100A1
UniProt Synonym Gene Names	S100A
UniProt Entry Name	S10A1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/s100-antibody-clone-jm51-31-western-blot-immunoprecipitation-immunohistochemistry-flow-cytometry-elisa/30331>