

Rabbit anti-Human, Mouse TSC2 Polyclonal Antibody | anti-TSC2 antibody

TSC2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA303614	Rabbit	Human, Mouse	Immunocytochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

TSC2; Polyclonal Antibody; TSC2 Antibody; Tuberous sclerosis complex 2; Tuberin; anti-TSC2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Purity/Purification	Affinity chromatography purified via peptide column
Form/Format	Supplied in PBS containing 0.02% sodium azide.
Sequence Length	1807
Applicable Applications for anti-TSC2 antibody	ICC (Immunocytochemistry), WB (Western Blot), ELISA
Immunogen Type	Peptide
Immunogen Description	TSC2 (NT) was raised against a 14 amino acid peptide from near the amino terminus of human TSC2.
Target Name	TSC2
Preparation and Storage	Can be stored at -20 degree C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

IMAGES



ICC (Immunocytochemistry)

Immunocytochemistry of TSC2 in L1210 cells with TSC2 antibody at 10 ug/mL.



WB (Western Blot)

Western blot analysis of TSC2 in L1210 cell lysate with TSC2 antibody at (A) 2 and (B) 4 ug/mL.

ADDITIONAL INFORMATION

Related Product Information for anti-TSC2 antibody	Tuberous sclerosis complex (TSC) is an autosomal dominant tumor syndrome caused by mutations in either of the TSC1 or TSC2 tumor suppressor genes. The products of these genes form a protein complex that indirectly decreases the signaling of the mammalian Target of Rapamycin (TOR), an evolutionarily conserved serine/threonine kinase that regulates cell growth and cell cycle through its ability to integrate signals from nutrient levels and growth factors. TOR activity is stimulated by Rheb, a member of the Ras superfamily of G-proteins, when the GTP/GDP ratio bound to Rheb is high. Immunoprecipitated TSC1/TSC2 has been shown to stimulate Rheb GTPase activity in vitro, suggesting that the TSC1/TSC2 complex decreases the ability of Rheb to stimulate TOR activity. This is supported by experiments showing that overexpression of TSC1 and TSC2 results in a significant decrease in the GTP/GDP ratio bound to Rheb and the inhibition of cell growth. At least three isoforms of TSC2 exist.
Product Categories/Family for anti-TSC2 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	116256352
NCBI GeneID	7249
NCBI Accession #	NP_000539
NCBI GenBank Nucleotide #	NM_000548.3
Molecular Weight	25,773 Da
NCBI Official Full Name	tuberin isoform 1
NCBI Official Synonym Full Names	tuberous sclerosis 2
NCBI Official Symbol	TSC2
NCBI Official Synonym Symbols	LAM; TSC4; PPP1R160
NCBI Protein Information	tuberin
UniProt Protein Name	Tuberin
UniProt Gene Name	TSC2
UniProt Synonym Gene Names	TSC4
UniProt Entry Name	TSC2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tsc2-antibody-immunocytochemistry-western-blot-elisa/303614>