

Rabbit FGF2 Polyclonal Antibody | anti-FGF2 antibody

FGF2 Antibody

CATALOG # AAA303903	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

FGF2; Polyclonal Antibody; FGF2 Antibody; BFGF; FGFB; FGF-2; HBGF-2; anti-FGF2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of total FGF2 protein.
Purity/Purification	Antigen affinity purification.
Form/Format	Rabbit IgG in pH7.3 PBS, 0.05% NaN ₃ , 50% Glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	288
Applicable Applications for anti-FGF2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide corresponding to a region derived from internal residues of human Fibroblast growth factor 2
Target Name	FGF2
Preparation and Storage	Store at -20 degree C

IMAGES


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human breast cancer tissue using at dilution 1/100.


Application Data

Gel: 8% SDS-PAGE Lysates (from left to right): Mouse lung tissue
Amount of lysate: 30ug per lane
Primary antibody: 1/1000 dilution
Secondary antibody dilution: 1/8000
Exposure time: 10 seconds

ADDITIONAL INFORMATION

Related Product Information for anti-FGF2 antibody	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF.
Product Categories/Family for anti-FGF2 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	153285461
NCBI GeneID	2247
NCBI Accession #	NP_001997
NCBI GenBank Nucleotide #	NM_002006.4
UniProt Accession #	P09038; O00527; P78443; Q16443; Q5PY50; Q7KZ11; Q7KZ72; Q9UC54; Q9UCS5; Q9UCS6; A4LBB8
Molecular Weight	SDS-PAGE MW: 31kd
NCBI Official Full Name	fibroblast growth factor 2
NCBI Official Synonym Full Names	fibroblast growth factor 2 (basic)
NCBI Official Symbol	FGF2
NCBI Official Synonym Symbols	BFGF; FGFB; FGF-2; HBGF-2
NCBI Protein Information	fibroblast growth factor 2
UniProt Protein Name	Fibroblast growth factor 2
UniProt Gene Name	FGF2
UniProt Synonym Gene Names	FGFB; FGF-2; bFGF; HBGF-2
UniProt Entry Name	FGF2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fgf2-antibody-immunohistochemistry-western-blot-elisa/303903>