

Rabbit Tau Polyclonal Antibody | anti-Tau antibody

Tau (Phospho-Ser262) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA304262	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Tau; Polyclonal Antibody; Tau (Phospho-Ser262) Antibody; MAPT; MTAPT; MTBT1; Neurofibrillary tangle protein; PHF-tau; anti-Tau antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of Tau only when phosphorylated at serine 262.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	776
Applicable Applications for anti-Tau antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 262 (I-G-S(p)-T-E) derived from Human Tau.
Modification	Phospho-Ser262
Target Name	Tau
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.


IMAGES



IF (Immunofluorescence)
Immunofluorescence staining of methanol-fixed Hela cells using Tau(Phospho-Ser262) Antibody.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded rat hippocampal region tissue from a model with Alzheimer



WB (Western Blot)
Western blot analysis of extracts from mouse brain tissue using Tau(Phospho-Ser262) Antibody and the same antibody preincubated with blocking peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-Tau antibody	Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization.
Product Categories/Family for anti-Tau antibody	Phospho-specific Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	294862258
NCBI GeneID	4137
NCBI Accession #	NP_001116538.2
NCBI GenBank Nucleotide #	NM_001123066.3
UniProt Accession #	P10636; P18518; Q14799; Q15549; Q15550; Q15551; Q1RMF6; Q53YB1; Q5CZI7; Q5XWF0; Q6QT54; Q9UDJ3
Molecular Weight	80,941 Da
NCBI Official Full Name	microtubule-associated protein tau isoform 6
NCBI Official Synonym Full Names	microtubule-associated protein tau
NCBI Official Symbol	MAPT

NCBI Official Synonym Symbols	TAU; MSTD; PPND; DDPAC; MAPTL; MTBT1; MTBT2; FTDP-17; PPP1R103
NCBI Protein Information	microtubule-associated protein tau
UniProt Protein Name	Microtubule-associated protein tau
UniProt Gene Name	MAPT
UniProt Synonym Gene Names	MAPTL; MTBT1; TAU; PHF-tau
UniProt Entry Name	TAU_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tau-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/304262>