

Rabbit PDIA3 Polyclonal Antibody | anti-PDIA3 antibody

PDIA3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA304962	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PDIA3; Polyclonal Antibody; PDIA3 Antibody; ER60; ERp57; ERp60; ERp61; anti-PDIA3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total PDIA3 protein.
Purity/Purification	Antibodies were purified by affinity purification using immunogen.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	505
Applicable Applications for anti-PDIA3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Recombinant Protein
Immunogen Description	Recombinant protein of human PDIA3.
Target Name	PDIA3
Preparation and Storage	Store at -20 degree C

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cell using PDIA3 antibody. Blue: DAPI for nuclear staining.



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded rat kidney using PDIA3 antibody at dilution of 1:200 (200x lens).



WB (Western Blot)
Western blot analysis of extracts of various cell lines using PDIA3 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-PDIA3 antibody	Secretory proteins translocate into the endoplasmic reticulum (ER) after their synthesis where they are post-translationally modified and properly folded. To reach their native conformation, many secretory proteins require the formation of intra- or inter-molecular disulfide bonds (1). This process is called oxidative protein folding. Disulfide isomerase (PDI) has two thioredoxin homology domains and catalyzes the formation and isomerization of these disulfide bonds (2). Other ER resident proteins that possess the thioredoxin homology domains, including endoplasmic reticulum stress protein 57 (ERp57), constitute the PDI family (2). ERp57 interacts with calnexin and calreticulin (3) and is suggested to play a role in the isomerization of disulfide bonds on certain glycoproteins (3).
Product Categories/Family for anti-PDIA3 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	21361657
NCBI GeneID	2923
NCBI Accession #	NP_005304.3
NCBI GenBank Nucleotide #	NM_005313.4
UniProt Accession #	P30101; Q13453; Q14255; Q8IYF8; Q9UMU7; AAB37397
Molecular Weight	56,782 Da
NCBI Official Full Name	protein disulfide-isomerase A3
NCBI Official Synonym Full Names	protein disulfide isomerase family A, member 3
NCBI Official Symbol	PDIA3
NCBI Official Synonym Symbols	P58; ER60; ERp57; ERp60; ERp61; GRP57; GRP58; PI-PLC; HsT17083; HEL-S-269; HEL-S-93n
NCBI Protein Information	protein disulfide-isomerase A3
UniProt Protein Name	Protein disulfide-isomerase A3
UniProt Gene Name	PDIA3
UniProt Synonym Gene Names	ERP57; ERP60; GRP58; p58; ER protein 57; ERp57; ER protein 60; ERp60
UniProt Entry Name	PDIA3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.