

Rabbit anti-Human WISP1 Polyclonal Antibody | anti-WISP1 antibody

WISP1 Antibody

CATALOG # AAA305282	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

WISP1; Polyclonal Antibody; WISP1 Antibody; CCN4; WISP1c; WISP1i; WISP1tc; anti-WISP1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of total WISP1 protein.
Purity/Purification	Antigen affinity purification.
Form/Format	Rabbit IgG in pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol.
Concentration	1.6 mg/ml (varies by lot)
Sequence Length	155
Applicable Applications for anti-WISP1 antibody	IHC (Immunohistochemistry)
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide of human WNT1 inducible signaling pathway protein 1
Target Name	WISP1
Preparation and Storage	Store at -20 degree C

IMAGES


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human thyroid cancer tissue using at dilution 1/25.


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human brain tissue using at dilution 1/25.

ADDITIONAL INFORMATION

Related Product Information for anti-WISP1 antibody	This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like domain. This gene may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. It is expressed at a high level in fibroblast cells, and overexpressed in colon tumors. The encoded protein binds to decorin and biglycan, two members of a family of small leucine-rich proteoglycans present in the extracellular matrix of connective tissue, and possibly prevents the inhibitory activity of decorin and biglycan in tumor cell proliferation. It also attenuates p53-mediated apoptosis in response to DNA damage through activation of the Akt kinase. It is 83% identical to the mouse protein at the amino acid level. Multiple alternatively spliced transcript variants have been identified.
Product Categories/Family for anti-WISP1 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	325910843
NCBI GeneID	8840
NCBI Accession #	NP_001191798.1
NCBI GenBank Nucleotide #	NM_001204869.1
UniProt Accession #	O95388; Q5JBS6; Q5JBS7; Q5JBS8; Q9HCS3; A8KAG6; E7EMM5
Molecular Weight	21,550 Da
NCBI Official Full Name	WNT1-inducible-signaling pathway protein 1 isoform 3
NCBI Official Synonym Full Names	WNT1 inducible signaling pathway protein 1
NCBI Official Symbol	WISP1
NCBI Official Synonym Symbols	CCN4; WISP1c; WISP1i; WISP1tc
NCBI Protein Information	WNT1-inducible-signaling pathway protein 1
UniProt Protein Name	WNT1-inducible-signaling pathway protein 1
UniProt Gene Name	WISP1
UniProt Synonym Gene Names	CCN4; WISP-1
UniProt Entry Name	WISP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/wisp1-antibody-immunohistochemistry-elisa/305282>