

Rabbit TLR2 Polyclonal Antibody | anti-TLR2 antibody

TLR2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA30547	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

TLR2; Polyclonal Antibody; TLR2 Polyclonal Antibody; CD282; TIL4; anti-TLR2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH7.4, 150mM NaCl, 0.02% sodium azide and 158% glycerol. PBS with 0.02% sodium azide, 50% glycerol, pH7.111
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-TLR2 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200 IF: 1:50-1:100
Immunogen	A synthetic peptide of human TLR2
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.


IMAGES


IF (Immunofluorescence)

Immunofluorescence analysis of mouse lung using TLR2 antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat spleen using TLR2 antibody at dilution of 1:100.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse islets of langerhans using TLR2 antibody at dilution of 1:150 .




WB (Western Blot)

Western blot analysis of extracts of various cell lines, using TLR2 antibody at 1:1000 dilution.



IF (Immunofluorescence)

Immunofluorescence analysis of rat lung using TLR2 antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cells using TLR2 antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

ADDITIONAL INFORMATION

Related Product Information for anti-TLR2 antibody	The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. This protein is a cell-surface protein that can form heterodimers with other TLR family members to recognize conserved molecules derived from microorganisms known as pathogen-associated molecular patterns (PAMPs). Activation of TLRs by PAMPs leads to an up-regulation of signaling pathways to modulate the host's inflammatory response. This gene is also thought to promote apoptosis in response to bacterial lipoproteins. This gene has been implicated in the pathogenesis of several autoimmune diseases. Alternative splicing results in multiple transcript variants.
Product Categories/Family for anti-TLR2 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	19718734
NCBI GeneID	7097
NCBI Accession #	NP_003255.2
NCBI GenBank Nucleotide #	NM_003264.3
UniProt Accession #	O60603; O15454; Q8NI00; B3Y612; D1CS45; D1CS48; D1CS49
Molecular Weight	89,838 Da
NCBI Official Full Name	toll-like receptor 2

NCBI Official Synonym Full Names	toll-like receptor 2
NCBI Official Symbol	TLR2
NCBI Official Synonym Symbols	TIL4; CD282
NCBI Protein Information	toll-like receptor 2; toll/interleukin 1 receptor-like 4; toll/interleukin-1 receptor-like protein 4
UniProt Protein Name	Toll-like receptor 2
UniProt Gene Name	TLR2
UniProt Synonym Gene Names	TIL4
UniProt Entry Name	TLR2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tlr2-antibody-western-blot-immunohistochemistry-immunofluorescence-elisa/30547>