

Rabbit anti-Human ACVR2B Polyclonal Antibody | anti-ACVR2B antibody

ACVR2B Antibody

CATALOG # AAA305568	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

ACVR2B; Polyclonal Antibody; ACVR2B Antibody; HTX4; ACTRIIB; ActR-IIB; anti-ACVR2B antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of total ACVR2B protein.
Purity/Purification	Antigen affinity purification.
Form/Format	Rabbit IgG in pH7.3 PBS, 0.05% NaN ₃ , 50% Glycerol.
Concentration	1.1 mg/ml (varies by lot)
Sequence Length	512
Applicable Applications for anti-ACVR2B antibody	IHC (Immunohistochemistry)
Immunogen Type	Recombinant Protein
Immunogen Description	Fusion protein corresponding to residues near the C terminal of human activin A receptor, type IIB
Target Name	ACVR2B
Preparation and Storage	Store at -20 degree C

IMAGES


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human cervical cancer tissue using at dilution 1/25.


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human thyroid cancer tissue using at dilution 1/25.

ADDITIONAL INFORMATION

Related Product Information for anti-ACVR2B antibody	Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. This gene encodes activin A type IIB receptor, which displays a 3- to 4-fold higher affinity for the ligand than activin A type II receptor.
Product Categories/Family for anti-ACVR2B antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	116734708
NCBI GeneID	93
NCBI Accession #	NP_001097.2
NCBI GenBank Nucleotide #	NM_001106.3
UniProt Accession #	Q13705; Q4VAV0
Molecular Weight	– Da
NCBI Official Full Name	activin receptor type-2B
NCBI Official Synonym Full Names	activin A receptor, type IIB
NCBI Official Symbol	ACVR2B
NCBI Official Synonym Symbols	HTX4; ACTRIIB; ActR-IIB
NCBI Protein Information	activin receptor type-2B
UniProt Protein Name	Activin receptor type-2B
UniProt Gene Name	ACVR2B
UniProt Synonym Gene Names	ACTR-IIB
UniProt Entry Name	AVR2B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/acvr2b-antibody-immunohistochemistry-elisa/305568>