

Rabbit anti-Human Factor X Polyclonal Antibody | anti-F10 antibody

Factor X antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA305579	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Factor X; Polyclonal Antibody; Factor X antibody; anti-F10 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Purified by antigen-affinity chromatography.
Form/Format	Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a preservative.
Sequence Length	488
Applicable Applications for anti-F10 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Recombinant protein
Immunogen Description	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 33 and 282 of Human F10
Target Name	Factor X
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of methanol-fixed HeLa, using Factor X antibody at 1: 200 dilution.



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded A549 xenograft, using Factor X antibody at 1: 500 dilution.



Application Data

Sample (30 ug of whole cell lysate)A: Hep G2 10% SDS PAGEPrimary antibody diluted at 1:5000

ADDITIONAL INFORMATION

Related Product Information for anti-F10 antibody	This gene encodes the vitamin K-dependent coagulation factor X of the blood coagulation cascade. This factor undergoes multiple processing steps before its preproprotein is converted to a mature two-chain form by the excision of the tripeptide RKR. Two chains of the factor are held together by 1 or more disulfide bonds; the light chain contains 2 EGF-like domains, while the heavy chain contains the catalytic domain which is structurally homologous to those of the other hemostatic serine proteases. The mature factor is activated by the cleavage of the activation peptide by factor IXa (in the intrinsic pathway), or by factor VIIa (in the extrinsic pathway). The activated factor then converts prothrombin to thrombin in the presence of factor Va, Ca+2, and phospholipid during blood clotting. Mutations of this gene result in factor X deficiency, a hemorrhagic condition of variable severity. [provided by RefSeq]
Product Categories/Family for anti-F10 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503625
NCBI GeneID	2159
NCBI Accession #	NP_000495
NCBI GenBank Nucleotide #	NM_000504.3
Molecular Weight	54,732 Da
NCBI Official Full Name	coagulation factor X isoform 1 preproprotein
NCBI Official Synonym Full Names	coagulation factor X
NCBI Official Symbol	F10
NCBI Official Synonym Symbols	FX; FXA
NCBI Protein Information	coagulation factor X
UniProt Protein Name	Coagulation factor X
UniProt Gene Name	F10
UniProt Entry Name	FA10_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.