

Rabbit p27 KIP 1 Polyclonal Antibody | anti-p27KIP1 antibody

p27 KIP 1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA30578	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence

PRODUCT SYNONYMS

p27 KIP 1; Polyclonal Antibody; p27 KIP 1 Polyclonal Antibody; CDKN1B; CDKN4; KIP1; MEN1B; MEN4; P27KIP1; anti-p27KIP1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH7.4, 150mM NaCl, 0.02% sodium azide and 108% glycerol. PBS with 0.02% sodium azide, 50% glycerol, pH7.61
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-p27KIP1 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:100 IF: 1:50-1:200
Immunogen	A synthetic peptide of human p27 KIP 1 (NP_004055.1)
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse testis using p27 KIP 1 antibody at dilution of 1:100 .




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human gastric cancer using p27 KIP 1 antibody at dilution of 1:100 .



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human breast using p27 KIP 1 antibody at dilution of 1:100 .



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon using p27 KIP 1 antibody at dilution of 1:100 .

**IHC (Immunohistochemistry)**

Immunohistochemistry of paraffin-embedded human tonsil using p27 KIP 1 antibody at dilution of 1:100 .

**IHC (Immunohistochemistry)**

Immunohistochemistry of paraffin-embedded rat spleen using p27 KIP 1 antibody at dilution of 1:100 .

ADDITIONAL INFORMATION

Related Product Information for anti-p27KIP1 antibody	This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. Mutations in this gene are associated with multiple endocrine neoplasia type IV (MEN4).
Product Categories/Family for anti-p27KIP1 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757962
NCBI GeneID	1027
NCBI Accession #	NP_004055.1
NCBI GenBank Nucleotide #	NM_004064.3
UniProt Accession #	P46527; Q16307; Q5U0H2; Q9BUS6
Molecular Weight	22,073 Da
NCBI Official Full Name	cyclin-dependent kinase inhibitor 1B
NCBI Official Synonym Full Names	cyclin-dependent kinase inhibitor 1B (p27, Kip1)
NCBI Official Symbol	CDKN1B
NCBI Official Synonym Symbols	KIP1; MEN4; CDKN4; MEN1B; P27KIP1
NCBI Protein Information	cyclin-dependent kinase inhibitor 1B
UniProt Protein Name	Cyclin-dependent kinase inhibitor 1B
UniProt Gene Name	CDKN1B
UniProt Synonym Gene Names	KIP1
UniProt Entry Name	CDN1B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/p27-kip-1-antibody-western-blot-immunohistochemistry-immunofluorescence-elisa/30578>