

Rabbit IL2 Polyclonal Antibody | anti-IL2 antibody

IL2 Rabbit Polyclonal Antibody

CATALOG # AAA30640	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

IL2; Polyclonal Antibody; IL2 Rabbit Polyclonal Antibody; IL-2; TCGF; lymphokine; anti-IL2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-IL2 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200
Immunogen	Recombinant protein of human IL2
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.


IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse spleen using IL2 at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse lung using IL2 at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human lung using IL2 at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human tonsil using IL2 at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of Raw264.7 cells, using IL2 at 1:1000 dilution. Raw264.7 cells were treated by LPS for 6 hours and Brefeldin A for 3 hours of stimulation.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using IL2 at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-IL2 antibody

The protein encoded by this gene is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli.

Product Categories/Family for anti-IL2 antibody

[Primary Antibody](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	28178861
NCBI GeneID	3558
NCBI Accession #	NP_000577.2
NCBI GenBank Nucleotide #	NM_000586.3
UniProt Accession #	P60568 ; P01585
Molecular Weight	17,628 Da
NCBI Official Full Name	interleukin-2
NCBI Official Synonym Full Names	interleukin 2

NCBI Official Symbol	IL2
NCBI Official Synonym Symbols	IL-2; TCGF; lymphokine
NCBI Protein Information	interleukin-2; aldesleukin; T cell growth factor; involved in regulation of T-cell clonal expansion
UniProt Protein Name	Interleukin-2
UniProt Gene Name	IL2
UniProt Synonym Gene Names	IL-2; TCGF
UniProt Entry Name	IL2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/il2-antibody-western-blot-immunohistochemistry-elisa/30640>